

Instructions for Patients

By completing this form, you can:



Learn about your health insurance coverage and financial assistance options through Genentech MySMA Support™



Sign up to receive **optional** disease education and other materials, including **optional** services from Genentech MySMA Support

You can choose not to sign this form. However, Genentech cannot provide you with your health insurance benefits investigation and other financial assistance options without your signed authorization on page 4. Enrollment in this program does not impact your ability to gain access to Evrysdi from your health care provider or health insurance plan.

▶ Please follow these steps to get started:

- 1 Read** "Authorization to Use and Disclose Personal Information" on page 3.
- 2 Sign and date** page 4. Please note you must sign the form to get support for your treatment.
- 3 Send** in your completed form using one of the options below.

Genentech can start supporting you when **page 4** of this form is submitted by you or your doctor's office in one of the following ways:



Complete online by scanning this QR code or visiting www.evrysdi.com/forms

OR



Print, complete, take a photo and text it to **(650) 877-1111**

OR



Print, complete and fax it to **(833) 387-9700**

Please write legibly and complete all required fields (*) on the Evrysdi Start Form to avoid any delays.

Please note: Your doctor has to complete the Evrysdi Prescriber Service Form before we can begin helping you.

If you have any questions, talk to your health care provider or call (833) 387-9734.

Helpful Terminology

Genentech: The maker of the medicine your doctor wants to prescribe. Genentech is committed to helping patients get the medicine their doctor prescribed. When used on this form, “Genentech” refers to Genentech, Genentech Patient Foundation, and their respective partners, affiliates, subcontractors and agents.

MySMA Support™: Your support team at Genentech that works with your doctor and your health insurance plan to help you understand your insurance coverage and get your prescribed Evrysdi medicine. The Genentech MySMA team includes your Case Manager (CM), specialty pharmacy, and a Partnership and Access Liaison (PAL).

Additional Partnership and Access Liaison (PAL) Support:

A local representative from Genentech that offers **optional** disease education and product support for patients at no cost to them. This may include items or materials explaining product dosing and administration for use when traveling and may also include marketing materials and information about Genentech products, services and programs. Please keep in mind that PALs are not part of your medical team, do not provide medical advice and are not substitutes for your health care provider. Your health care provider should always be your main resource for any questions about your health and medical care.

Case Manager (CM): The Genentech representative that partners closely with your health care provider, and other members of the MySMA Support team, to help you understand your health insurance coverage and potential financial support options for Evrysdi.

Specialty pharmacy (SP): An SP supplies certain medicines for patients. Some plans require you to use a certain SP to receive your medicine. SPs send your medicine to your doctor’s office or your home. They

may also offer other services, such as referrals to financial assistance.

Genentech Patient Foundation: A program that gives free Genentech medicine to people who don’t have health insurance coverage or who have financial concerns and meet certain eligibility criteria.

Household size: Number of people living in your household, including you.

Net household income: How much you and the members of your household currently make each year minus specific deductions. This is also frequently referred to as your Adjusted Gross Income or AGI. This information is needed to determine Genentech Patient Foundation eligibility.

Deductible: The amount you pay for your health care services or medicines out of pocket before your health insurance plan begins to pay.

Out-of-pocket costs: The amount not paid by the health insurance plan that you must pay for your treatment. This includes premiums, deductibles, co-pays and co-insurance.

Co-pay assistance: Programs available to help eligible patients pay for their medicines.

Alternate contact: Someone you choose to be your contact person if Genentech MySMA Support cannot reach you. An Alternate Contact may not be an individual associated with or a representative of your insurance company, employer, or a business partner of your insurance company or employer.

Legally authorized representative: An individual or judicial or other body authorized under applicable law to consent on behalf of a patient (e.g., parent or legal guardian of a minor).

Terms and Conditions of the Genentech Patient Foundation

- If I receive free medicine from the Genentech Patient Foundation, I will not sell or give out the medicine because it is illegal to do so. I am responsible to ensure that the medicine is sent to a secure address when shipped to me, and I must control any medicine that I receive
- I understand that, for purposes of an audit, the Genentech Patient Foundation could ask me for a copy of my IRS 1040 form or other proof of income
- Some insurance plans and/or employers partner with organizations known as alternate funding programs. Such arrangements require patients to apply to the Genentech Patient Foundation as a condition of, or prerequisite to, coverage of relevant Genentech products. These alternate funding programs include SHARx, Paydhealth, and Payer Matrix, among others. Patients whose insurance plans and/or employers use an alternative funding program are ineligible for support from the Genentech Patient Foundation
- I acknowledge that, to the best of my knowledge, neither my insurance plan nor my employer (1) required me to apply to the Genentech Patient Foundation and/or (2) changed or hid my insurance coverage for my Genentech medicine to make me appear to be underinsured and eligible for support from the Genentech Patient Foundation. I am not applying to the Genentech Patient Foundation on behalf of someone whose insurance plan and/or employer partners with an alternative funding program. The Alternate Contact listed on my application (if any) is not associated with or a representative of my insurance company, employer, or a business partner of my insurance company or employer. If I subsequently learn that my insurance plan and/or employer uses an alternative funding program, I agree to inform the Genentech Patient Foundation immediately and understand that I will no longer be eligible for support

Authorization to Use and Disclose Personal Information

I authorize my physician(s) and their staff, pharmacies, and health insurance plan (my “health care providers”) to share my personal information, which may include contact information, demographic information, financial information, and information related to my medical condition, treatments, and health insurance and benefits, with Genentech, Genentech Patient Foundation, and their respective partners, affiliates, subcontractors, and agents (together, “Genentech”). I authorize Genentech to receive, use, and share my personal information in order to provide me with access to the products, services, and programs described on this form, which may include the following:

- Working with my health insurance plan to understand or verify coverage for Genentech products
- Applying to the Genentech Patient Foundation
- Determining my eligibility for and facilitating enrollment into financial assistance services if I’m eligible, including co-pay assistance
- Coordinating my prescription through a pharmacy, infusion site and/or health care provider’s office. This includes contacting me to discuss my coverage, costs and eligibility for assistance and other program administration purposes
- Facilitating my access to Genentech products
- Ensuring quality and safety and improving our products and services
- Contacting me by mail, e-mail, telephone calls and text messages at the number(s) and address(es) provided for non-marketing purposes
- If I agree to the **optional** Consent for Patient Resources and Information, providing me with **optional** disease information and marketing material about products, services and programs offered by Genentech, its partners and their respective affiliates. This includes **optional** services or engagement from Genentech MySMA Support, which may include outreach by a PAL. This is not required to receive help from Genentech MySMA Support with understanding health insurance coverage and potential financial support programs
- If I agree to opt into marketing autodialed and texted communications, contacting me by autodialed calls and/or text messages at the phone number(s) I have provided for marketing purposes, including from a PAL. This is not required to receive help from Genentech MySMA Support with understanding health insurance coverage and potential financial support programs

I understand that this will include sharing and use of information about me that could be considered sensitive personal information, such as health conditions, but that the use of this information by Genentech is necessary to determine if I qualify for and to administer the benefits and services for which I am applying. I understand that Genentech may also share my personal information, including sensitive personal information, for the purposes described on this authorization with my health care providers, service providers, and any individual I may designate as an alternate contact. I understand that my pharmacy may receive payment or other remuneration for disclosing my personal information pursuant to this authorization. I can choose not to sign this authorization, but Genentech will not be able to provide the services to me without it. However, my health care providers may not condition either my treatment or my payment, enrollment, or eligibility for benefits on signing this authorization.

I also understand and agree that:

- This authorization is valid for 6 years from the date I sign or the date I last enrolled, whichever comes first, unless a shorter period is required by law, or I revoke it earlier
- My personal information released under this authorization may no longer be protected by state and federal law, including the Health Insurance Portability and Accountability Act (HIPAA). However, Genentech will only use and share my personal information for the purposes stated on this authorization or as otherwise permitted by law
- I have the right to revoke (cancel) this authorization at any time by submitting a written notice to: Genentech Access Solutions, 1 DNA Way, South San Francisco, CA 94080-4990 or by calling **(866) 422-2377**. If I revoke this authorization, I will no longer be eligible for the services described. If a health care provider is disclosing my personal information to Genentech on an authorized, ongoing basis, my revocation will be effective with respect to such health care provider when they receive notice of my revocation. My revocation will not impact uses and disclosures of my personal information that have already occurred in reliance on this authorization
- More information on my privacy rights, including specific rights I may have as a resident of certain states, can be found in Genentech’s **Privacy Policy** (www.gene.com/privacy-policy)
- I have a right to receive a copy of this authorization

Patient Information (to be completed by patient or their legally authorized representative)

*First name: _____ *Last name: _____

Home phone: _____ Cell phone: _____

OK to leave a detailed message? Date of birth (MM/DD/YYYY): _____ / _____ / _____

Email: _____ Preferred language: English Spanish Other: _____

Alternate Contact (optional) Full name: _____

Relationship: _____ Phone: _____

1 Financial Eligibility: Complete **only** if you are applying to the Genentech Patient Foundation By completing this section, I am agreeing to the Terms and Conditions of the Genentech Patient Foundation outlined on page 2.
Household size (including you): _____
Annual household income: _____

2 Consent for Patient Resources and Information (OPTIONAL)
Genentech offers disease education and product support for patients, including items or marketing materials explaining the product and how to take it, use when traveling with the product and other information about Genentech products, services and programs. You do not have to sign up for these resources and support to get help with your insurance coverage or to learn about financial assistance options. Signing up here allows you to be contacted using the information you provide on this form. These marketing materials and support are **optional**, free and may be provided by a PAL, Genentech’s partners and their respective affiliates. PALs do not provide medical advice. Your healthcare provider should always be your main resource for any questions about your health and medical care.

By checking this box, I agree to receive disease education materials and product support services, including outreach by a PAL. I understand that I don’t have to opt into this offer and my decision does not affect receiving my medicine or financial support information. It may be necessary to use my sensitive personal information to provide me with relevant material. I also understand that I may opt out of receiving this information at any time by calling **(877) 436-3683**.

By checking this box, I agree to receive autodialed calls and text messages, which may include marketing communications about Evrysdi from and on behalf of Genentech, including from a PAL, at the phone number(s) provided. I understand that choosing to receive these messages is voluntary and is not a requirement of any purchase or program enrollment. Message frequency may vary. Message and data rates may apply. I may opt out at any time by texting STOP or calling **(877) GENENTECH/(877) 436-3683**. I am also agreeing to the **Privacy Policy (www.gene.com/privacy-policy)** and **SMS Terms & Conditions (www.gene.com/terms-conditions/sms-text-message-program-terms-conditions)**.

3 By signing this form, I acknowledge that I have provided accurate and complete information and understand and agree to the terms of this form. My signature certifies that I have read, understood, and agree to the release and use of my personal information, including sensitive personal information, pursuant to the Authorization to Use and Disclose Personal Information and as otherwise stated on this form.

REQUIRED	Sign and date here	_____ / _____ / _____
		*Signature of Patient/Legally Authorized Representative (A parent or guardian must sign for patients under 18 years of age)
		*Date signed (MM/DD/YYYY)
	Person signing (if not patient)	_____
	Print first name	_____
	Print last name	_____
	Relationship to patient	_____

Once this page (4/6) has been completed, please text a photo of the page to **(650) 877-1111** or fax to **(833) 387-9700**. You can also complete this form online at **www.evrysdi.com/forms**.

If this is an electronic consent, you understand that by typing your name and the date above and submitting, or taking a picture and sending to us, that you are providing your consent electronically and that it has the same force and effect as if you were signing in person on paper. Genentech reserves the right to rescind, revoke or amend the program without notice at any time.

Instructions for Health Care Providers

By completing this form, you are requesting services on behalf of your patient, which may include:



Insurance benefits investigation



Resources for prior authorizations and appeals



Referrals of eligible patients to co-pay support options or the Genentech Patient Foundation

▶ To enroll your patient, please follow these steps:

- 1** Have your patient read pages 1-3.
- 2** Have your **patient complete the Patient Information on page 4** and sign and date Section 3:
 - Only the Patient Information and Section 3 are required for insurance coverage and financial assistance options support
 - If your patient is requesting free medicine from the Genentech Patient Foundation, they should also complete Section 1
 - If your patient is requesting **optional** disease education and other material, including **optional** services from Genentech MySMA Support™, they should also complete Section 2
- 3** **Complete page 6 and sign and date** Section 7.
- 4** **Submit pages 4 and 6 of the Start Form** via fax to **(833) 387-9700** or eSubmit at **www.evrysdi.com/forms**. Page 4 of the Start Form can also be submitted by text to **(650) 877-1111** as indicated on page 1.

Please write legibly and complete all **required fields (*) on the Evrysdi Start Form to avoid any delays.**

Prescriber Service Form (to be completed by the prescriber)

Step 1 Patient Information

*First name: _____ *Last name: _____ Gender: Male Female
 *Date of birth (MM/DD/YYYY): ____ / ____ / ____ Preferred language: English Spanish Other: _____
 Street: _____ Apt: _____ City: _____ *State: _____ ZIP: _____
 Home phone: _____ Cell phone: _____ Do not contact patient
 Alternate contact name: _____ Relationship: _____ Alt. phone: _____

Step 2 Insurance Information

Is the patient insured? Yes No
 If the patient is a newborn, is the insurance policyholder attesting that the newborn has been or will be added to the insurance(s) listed below within the timeframe required by the insurance policy? (Note: Many insurers require addition within 30 days of birth) Yes No

If patient is uninsured, please refer to the Genentech Patient Foundation.

Please fill out the information below or attach a copy of the patient's medical and prescription insurance cards.

Primary Insurance	Secondary Insurance	Pharmacy Benefit
Insurance name		
Subscriber name (if not patient)		
Subscriber/Policy ID #		
Group #		
Insurance phone		

Step 3 Evrysdi Start Program (Signature Required)

Dispense 1-shipment supply: Oral solution _____ mg (_____ mL) once daily **OR** 5 mg once daily
 5-mg tablet
 1-time refill of option selected above; weight-based dosing requires a new Rx.
 Your signature authorizes the specialty pharmacy to dispense needed ancillary supplies for enteral administration of this medication, such as: ENFit® adapters, oral syringes, cassettes, administration sets and tubing.

For full eligibility criteria and Terms and Conditions, please visit www.genentech-pro.com/starter or speak to your Evrysdi representative. Genentech reserves the right to rescind, revoke, or amend the Programs without notice at any time.

Step 4 Diagnosis and Clinical Information

*Diagnosis code(s): G12.0 Infantile spinal muscular atrophy, type I G12.1 Other inherited spinal muscular atrophy
 G12.9 Spinal muscular atrophy, unspecified Other: _____
 SMA type: 0 1 2 3 4 SMN2 copy number: _____ Patient weight: _____ lbs kgs Date measured ____ / ____ / ____
 Has patient taken Evrysdi? Yes No Expected Evrysdi treatment start date: ____ / ____ / ____
 Previous therapy: Spinraza® (nusinersen) last dose: ____ / ____ / ____ Zolgensma® (onasemnogene abeparvovec-xioi) last dose: ____ / ____ / ____
 Other: _____ last dose: ____ / ____ / ____ Drug and non-drug allergies _____ No known allergies

Step 5 Prescription Information

Solution/Strength	Directions	Route	Quantity	Refills
<input type="checkbox"/> .75 mg/mL 80 mL	<input type="checkbox"/> Oral solution _____ mg (_____ mL) OR	<input type="checkbox"/> Oral <input type="checkbox"/> Feeding tube	<input type="checkbox"/> 1-month supply	
<input type="checkbox"/> 5-mg tablet	<input type="checkbox"/> 5 mg once daily <input type="checkbox"/> SIG: _____	Type: _____	<input type="checkbox"/> Other: _____	

Step 6 Prescriber Information

*First name: _____ *Last name: _____ *Practice name: _____
 *Street: _____ Suite: _____ *City: _____ *State: _____ *ZIP: _____
 Prescriber tax ID #: _____ Prescriber NPI[†] #: _____ Group NPI[†] #: _____
 Office contact: _____ Contact phone: _____ Contact fax: _____

If you are a resident of a US state that provides certain rights with respect to your personal information, a complete description of the personal information we may collect and process, the purposes for which it is used by Genentech, and your rights under your state's privacy laws concerning your personal information can be found in our privacy notice at www.gene.com/privacy-policy.

Step 7 Health Care Provider Certification

By submitting this form, I certify: (a) The above therapy is medically necessary for this patient and the treatment decision has been made by the prescribing physician; (b) If the indication for which I am prescribing a Genentech product is not listed in the FDA-approved label, I am prescribing the medication for an "unapproved" use, meaning that the FDA has not approved the efficacy, dosage amount or safety of this medication for such a use; (c) I received the authorization to release the information above and other protected health information (as defined by the Health Insurance Portability and Accountability Act of 1996 (HIPAA)) to Genentech, Inc., Genentech Access Solutions, the contracted dispensing pharmacy, or other contractors for the purpose of requesting reimbursement support, assisting in initiating or continuing therapy, as a break in treatment would negatively impact the patient's therapeutic outcome; (d) My patient meets the criteria for the Genentech Patient Foundation and to the best of my knowledge, this patient has no prescription insurance coverage (including Medicaid, Medicare, or other public or private programs) for the Genentech medicine listed above, or is unable to afford the cost-sharing requirements associated with his/her insurance coverage for this medication. If the patient is enrolled in an insurance plan, the plan does not require the patient's application to the Genentech Patient Foundation and/or has not changed or hidden the patient's coverage for the Genentech medicine to make them appear to be underinsured and eligible for the Genentech Patient Foundation; (e) The services I am requesting on behalf of the patient may include benefits investigation (BI), prior authorization support (PA), co-pay card and co-pay assistance foundation referral; (f) No action on these services will be taken until the patient consent document has been received; (g) I must comply with all state-specific prescription requirements, such as e-prescribing, state-specific prescription form, fax language, etc; I understand that noncompliance with state-specific requirements could result in outreach to me; (h) My patient meets the criteria for Genentech Patient Foundation (GPF); (i) I understand that Genentech reserves the right to modify or discontinue the program at any time and to verify the accuracy of information submitted; (j) I understand that the GPF does not provide free drug in the instance of an administrative error or a coverage restriction, such as a step edit. For certain products where the step edit may not be medically appropriate, as confirmed by the prescribing physician, the GPF may consider support following 1 level of appeal.



Sign, date & fax to
(833) 387-9700

_____/_____/_____
***Prescriber Signature — Dispense as Written** *Date *Prescriber Signature — Generic Substitution Permitted *Date
 (Original signature required) (Original signature required)

[†]National Provider Identifier.

All trademarks and trade names are the property of their respective owners.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use EVRYSDI safely and effectively. See full prescribing information for EVRYSDI.

EVRYSDI® (risdiplam) for oral solution

EVRYSDI® (risdiplam) tablets, for oral use

Initial U.S. Approval: 2020

RECENT MAJOR CHANGES

Dosage and Administration (2.1, 2.2)

2/2026

INDICATIONS AND USAGE

EVRYSDI is a survival of motor neuron 2 (SMN2) splicing modifier indicated for the treatment of spinal muscular atrophy (SMA) in pediatric and adult patients. (1)

DOSAGE AND ADMINISTRATION

- Administer once daily with or without food per the table below (2.1):

Age and Body Weight	Recommended Daily Dosage	Dosage Form
Less than 2 months of age	0.15 mg/kg	EVRYSDI for Oral Solution
2 months to less than 2 years of age	0.2 mg/kg	
2 years of age and older weighing less than 20 kg	0.25 mg/kg	
2 years of age and older weighing 20 kg or more	5 mg	EVRYSDI for Oral Solution or EVRYSDI Tablet

- Swallow EVRYSDI tablet whole with water or dispersed in non-chlorinated drinking water (e.g., filtered water). (2.2)
- Administer EVRYSDI for oral solution with the provided oral syringe. (2.2)

- EVRYSDI for oral solution must be constituted by a healthcare provider prior to dispensing. (2.4)

- See Full Prescribing Information for important preparation and administration instructions. (2.2, 2.4)

DOSAGE FORMS AND STRENGTHS

- For Oral Solution: 60 mg of risdiplam as a powder for constitution to provide 0.75 mg/mL solution. (3)
- Tablets: 5 mg

CONTRAINDICATIONS

None. (4)

ADVERSE REACTIONS

The most common adverse reactions in later-onset SMA (incidence at least 10% of patients treated with EVRYSDI and more frequent than control) were fever, diarrhea, and rash. (6.1)

The most common adverse reactions in infantile-onset SMA were similar to those observed in later-onset SMA patients. Additionally, adverse reactions with an incidence of at least 10% were upper respiratory tract infection, lower respiratory tract infection, constipation, vomiting, and cough. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Genentech at 1-888-835-2555 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

Avoid coadministration with drugs that are substrates of multidrug and toxin extrusion (MATE) transporters. (7.1)

USE IN SPECIFIC POPULATIONS

Pregnancy: Based on animal data, may cause fetal harm. (8.1)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 2/2026

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Dosing Information
- 2.2 Important Administration Instructions
- 2.3 Missed Dose
- 2.4 Preparation of Powder for Oral Solution by Healthcare Provider

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience

7 DRUG INTERACTIONS

- 7.1 Effect of EVRYSDI on Substrates of Multidrug and Toxin Extrusion (MATE) Protein Transporters

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 13.2 Animal Toxicology and/or Pharmacology

14 CLINICAL STUDIES

- 14.1 Infantile-Onset SMA
- 14.2 Later-Onset SMA
- 14.3 Pre-Symptomatic SMA

16 HOW SUPPLIED/STORAGE AND HANDLING

- 16.1 EVRYSDI for Oral Solution
- 16.2 EVRYSDI Tablets

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

EVERYSOI is indicated for the treatment of spinal muscular atrophy (SMA) in pediatric and adult patients.

2 DOSAGE AND ADMINISTRATION

2.1 Dosing Information

EVERYSOI is administered orally once daily with or without food at approximately the same time each day. The recommended dosage is determined by age and body weight (see Table 1). EVERYSOI tablets are available for patients prescribed the 5 mg dose.

Table 1 Adult and Pediatric Dosing Regimen by Age and Body Weight

Age and Body Weight	Recommended Daily Dosage	Dosage Form
Less than 2 months of age	0.15 mg/kg	EVERYSOI for Oral Solution
2 months to less than 2 years of age	0.2 mg/kg	
2 years of age and older weighing less than 20 kg	0.25 mg/kg	
2 years of age and older weighing 20 kg or more	5 mg	EVERYSOI for Oral Solution or EVERYSOI Tablet

2.2 Important Administration Instructions

It is recommended that a healthcare provider discuss with the patient or caregiver how to prepare the prescribed daily dose prior to administration of the first dose [see *Instructions for Use for EVERYSOI for Oral Solution and EVERYSOI Tablets and Patient Information*].

EVERYSOI for Oral Solution

In infants who are breastfed, EVERYSOI for oral solution can be administered before or after breastfeeding. EVERYSOI cannot be mixed with formula or milk.

Instruct patients or caregivers to administer the dose using the reusable oral syringe provided.

EVERYSOI for oral solution must be taken immediately after it is drawn up into the oral syringe. If EVERYSOI is not taken within 5 minutes, EVERYSOI should be discarded from the oral syringe, and a new dose should be prepared.

Instruct patients to drink water after taking EVERYSOI for oral solution to ensure the drug has been completely swallowed.

EVERYSOI for oral solution can be administered via a nasogastric or gastrostomy tube. The tube should be flushed with water after delivering EVERYSOI for oral solution [see *Instructions for Use*].

EVERYSOI Tablets

Swallow EVERYSOI tablets whole with water. Do not chew, cut, or crush the tablets.

Alternatively, the EVERYSOI tablet can also be dispersed in one teaspoon (5 mL) of room temperature non-chlorinated drinking water (e.g., filtered water). EVERYSOI tablets must not be dispersed in any liquid other than non-chlorinated drinking water. Do not expose the prepared dispersion to sunlight. Swirl the small cup gently for up to 3 minutes until fully mixed (though some particles will remain). Administer the dispersed tablet immediately. To ensure no particles

are left in the small cup, refill it with at least one tablespoon (15 mL) of non-chlorinated drinking water, swirl, and administer immediately again.

EVERYSOI must be taken immediately after it is dispersed in non-chlorinated drinking water. Discard the prepared dispersion if it is not used within 10 minutes of adding non-chlorinated drinking water.

The dispersed EVRYSDI tablet can be administered via a nasogastric or gastrostomy tube that is 8 French or higher. Flush the tube with the non-chlorinated drinking water [at least one tablespoon (15 mL)] used to rinse the dispersion cup [*see Instructions for Use*].

2.3 Missed Dose

If a dose of EVRYSDI is missed, EVRYSDI should be administered as soon as possible if still within 6 hours of the missed dose, and the usual dosing schedule can be resumed on the next day. Otherwise, the missed dose should be skipped, and the next dose should be taken at the regularly scheduled time on the next day.

If a dose is not fully swallowed or vomiting occurs after taking a dose of EVRYSDI, another dose should not be administered to make up for the lost dose. The patient should wait until the next day to take the next dose at the regularly scheduled time.

2.4 Preparation of Powder for Oral Solution by Healthcare Provider

EVRYSDI powder must be constituted to the oral solution by a pharmacist or other healthcare provider prior to dispensing to the patient.

Preparation of the EVRYSDI Oral Solution 0.75 mg/mL

The EVRYSDI “Instructions for Constitution” booklet contains more detailed instructions on the preparation of the oral solution [*see Instructions for Constitution*].

Caution should be exercised when handling EVRYSDI powder for oral solution. Avoid inhalation and direct contact with skin or mucous membranes with the dry powder and the constituted solution. If such contact occurs, wash thoroughly with soap and water; rinse eyes with water. Wear disposable gloves during the preparation and cleanup procedure.

1. Gently tap the bottom of the closed glass bottle to loosen the powder.
2. Remove the cap. Do not throw away the cap.
3. Carefully pour 79 mL of purified water into the EVRYSDI bottle to yield the 0.75 mg/mL oral solution. Do not mix EVRYSDI with formula or milk.
4. Insert the press-in bottle adapter into the bottle opening by pushing it down against the bottle lip. Ensure it is completely pressed against the bottle lip.
5. Re-cap the bottle tightly and shake well for 15 seconds. Wait for 10 minutes. You should have obtained a clear solution. If not, shake well again for another 15 seconds or until you have obtained a clear solution.
6. Write the date of expiration of the constituted oral solution (calculated as 64 days after constitution) and the lot number on the bottle label. Peel off the part of the bottle label that has the expiration date of the powder.
7. Put the bottle back in its original carton.
8. Select the appropriate oral syringes (1 mL, 6 mL, or 12 mL) based on the patient’s dosage and remove the other oral syringes from the carton.
9. Dispense with the “Instructions for Use” and FDA-approved patient labeling. Alert patients to read the important handling information described in the Instructions for Use.

Storage

Keep the constituted oral solution of EVRYSDI in the original amber bottle to protect from light. Store in a refrigerator at 2°C to 8°C (36°F to 46°F). Do not freeze. Discard any unused portion 64 days after constitution. Keep the bottle in an upright position with the cap tightly closed. If refrigeration is not available, EVRYSDI can be kept at room temperature up to 40°C (up to 104°F) for a combined total of 5 days. EVRYSDI can be removed from, and returned to, a refrigerator. The total combined time out of refrigeration should not exceed 5 days.

3 DOSAGE FORMS AND STRENGTHS

EVRYSDI for oral solution: 60 mg as a light yellow, pale yellow, yellow, greyish yellow, greenish yellow, or light green powder for constitution. Following constitution, the volume of the greenish yellow to yellow solution is 80 mL, providing 60 mg/80 mL (0.75 mg/mL) risdiplam.

EVRYSDI tablet: 5 mg as a pale yellow film-coated tablet, round and curved, with EVR debossed on one side.

4 CONTRAINDICATIONS

None.

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in clinical trials of another drug and may not reflect the rates observed in practice.

In clinical trials including patients with infantile-onset SMA, later-onset SMA, and pre-symptomatic SMA, a total of 491 patients (51% female, 74% Caucasian) were exposed to EVRYSDI for up to a median duration of 48.1 months (range: 0.6 to 63.4 months), with 231 patients receiving treatment for more than 24 months. At the time of first EVRYSDI dose, 90 (18%) patients were 18 years and older, 119 (24%) were 12 years to less than 18 years, 189 (39%) were 2 years to less than 12 years, 67 (14%) 2 months to less than 2 years, and 26 (5%) were less than 2 months.

Clinical Trial in Later-Onset SMA

The safety of EVRYSDI for later-onset SMA is based on data from a randomized, double-blinded, placebo-controlled study (Study 2 Part 2) in patients with SMA Type 2 or 3 (n = 180) [see *Clinical Studies (14.2)*]. The patient population in Study 2 Part 2 ranged in age from 2 to 25 years at the time of the first dose.

The most common adverse reactions (reported in at least 10% of patients treated with EVRYSDI and at an incidence greater than on placebo) in Study 2 Part 2 were fever, diarrhea, and rash. Table 2 lists the adverse reactions that occurred in at least 5% of patients treated with EVRYSDI and at an incidence \geq 5% greater than on placebo in Study 2 Part 2.

Table 2 Adverse Reactions Reported in $\geq 5\%$ of Patients Treated with EVRYSDI and with an Incidence $\geq 5\%$ Greater Than on Placebo in Study 2 Part 2

Adverse Reaction	EVRYSDI (N = 120) %	Placebo (N = 60) %
Fever ¹	22	17
Diarrhea	17	8
Rash ²	17	2
Mouth and aphthous ulcers	7	0
Arthralgia	5	0
Urinary tract infection ³	5	0

¹ Includes pyrexia and hyperpyrexia.

² Includes rash, erythema, rash maculo-papular, rash erythematous, rash papular, dermatitis allergic, and folliculitis.

³ Includes urinary tract infection and cystitis.

Clinical Trial in Infantile-Onset SMA

The safety of EVRYSDI therapy for infantile-onset SMA is based on data from an open-label study in 62 patients (Study 1) [see *Clinical Studies (14.1)*]. The patient population ranged in age from 2 to 7 months at the time of the first EVRYSDI dose (weight range: 4.1 to 10.6 kg).

The most frequent adverse reactions reported in infantile-onset SMA patients treated with EVRYSDI in Study 1 were similar to those observed in later-onset SMA patients in Study 2. Additionally, the following adverse reactions reported in $\geq 10\%$ of patients were: upper respiratory tract infection (including nasopharyngitis, rhinitis), lower respiratory tract infection (including pneumonia, bronchitis), constipation, vomiting, and cough.

Clinical Trial in Pre-Symptomatic SMA

The safety of EVRYSDI therapy for pre-symptomatic SMA is based on data from an open-label, single-arm study in 26 patients (Study 3) [see *Clinical Studies (14.3)*]. The patient population ranged in age from 16 to 41 days at the time of the first dose (weight range: 3.1 to 5.7 kg). The safety profile of EVRYSDI in pre-symptomatic patients in Study 3 is consistent with the safety profile for symptomatic SMA patients treated with EVRYSDI in clinical trials.

7 DRUG INTERACTIONS

7.1 Effect of EVRYSDI on Substrates of Multidrug and Toxin Extrusion (MATE) Protein Transporters

Based on in vitro data, EVRYSDI may increase plasma concentrations of drugs eliminated via MATE1 or MATE2-K [see *Clinical Pharmacology (12.3)*], such as metformin. Avoid coadministration of EVRYSDI with MATE substrates. If coadministration cannot be avoided, monitor for drug-related toxicities and consider dosage reduction of the coadministered drug (based on the labeling of that drug) if needed.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Exposure Registry

There is a pregnancy exposure registry that monitors pregnancy and fetal/neonatal/infant outcomes in women exposed to EVRYSDI during pregnancy. Physicians are encouraged to

register patients and pregnant women are encouraged to register themselves by calling 1-833-760-1098 or visiting <https://www.evrysdipregnancyregistry.com>.

Risk Summary

There are no adequate data on the developmental risk associated with the use of EVRYSDI in pregnant women. In animal studies, administration of risdiplam during pregnancy or throughout pregnancy and lactation resulted in adverse effects on development (embryofetal mortality, malformations, decreased fetal body weights, and reproductive impairment in offspring) at or above clinically relevant drug exposures [see Data].

The estimated background risk of major birth defects and miscarriage for the indicated populations is unknown. In the U.S. general population, the estimated background risk of major birth defect and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively. Based on animal data, advise pregnant women of the potential risk to the fetus.

Data

Animal Data

Oral administration of risdiplam (0, 1, 3, or 7.5 mg/kg/day) to pregnant rats throughout organogenesis resulted in decreased fetal body weights and increased incidences of fetal structural variations at the highest dose tested, which was not associated with maternal toxicity. The no-effect level for adverse effects on embryofetal development (3 mg/kg/day) was associated with maternal plasma exposure (AUC) approximately 2 times that in humans at the maximum recommended human dose (MRHD) of 5 mg.

Oral administration of risdiplam (0, 1, 4, or 12 mg/kg/day) to pregnant rabbits throughout organogenesis resulted in embryofetal mortality, fetal malformations (hydrocephaly), and structural variations at the highest dose tested, which was associated with maternal toxicity. The no-effect dose for adverse effects on embryofetal development (4 mg/kg/day) was associated with maternal plasma exposure (AUC) approximately 4 times that in humans at the MRHD.

When risdiplam (0, 0.75, 1.5, or 3 mg/kg/day) was orally administered to rats throughout pregnancy and lactation, gestation was prolonged in the dams, and delayed sexual maturation (vaginal opening) and impaired reproductive function (decreased numbers of corpora lutea, implantation sites, and live embryos) were observed in female offspring at the highest dose. The no-effect dose for adverse effects on pre- and postnatal development in rats (1.5 mg/kg/day) was associated with maternal plasma exposure (AUC) similar to that in humans at the MRHD.

8.2 Lactation

Risk Summary

There are no data on the presence of risdiplam in human milk, the effects on the breastfed infant, or the effects on milk production. Risdiplam was excreted in the milk of lactating rats orally administered risdiplam.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for EVRYSDI and any potential adverse effects on the breastfed infant from EVRYSDI or from the underlying maternal condition.

8.3 Females and Males of Reproductive Potential

Studies of risdiplam in juvenile and adult rats and in monkeys demonstrated adverse effects on the reproductive organs, including germ cells, in males at clinically relevant plasma exposures [see *Use in Specific Populations (8.4) and Nonclinical Toxicology (13.1)*].

Pregnancy Testing

Pregnancy testing is recommended for females of reproductive potential prior to initiating EVRYSDI [see *Use in Specific Populations (8.1)*].

Contraception

EVRYSDI may cause embryofetal harm when administered to a pregnant woman [see *Use in Specific Populations (8.1)*].

Female Patients

Advise female patients of reproductive potential to use effective contraception during treatment with EVRYSDI and for at least 1 month after her last dose.

Infertility

Male Patients

Male fertility may be compromised by treatment with EVRYSDI [see *Nonclinical Toxicology (13.1)*].

Counsel male patients of reproductive potential receiving EVRYSDI about the potential effects on fertility. Male patients may consider sperm preservation prior to treatment.

8.4 Pediatric Use

The safety and effectiveness of EVRYSDI in pediatric patients (neonates and older) have been established. Use of EVRYSDI for SMA is supported by evidence from adequate and well-controlled studies of EVRYSDI in patients 2 months of age and older with SMA. Use of EVRYSDI for SMA in patients 2 months of age and younger is supported by pharmacokinetic and safety data from pediatric patients 16 days and older, and pharmacokinetic modeling and simulation to identify the dosing regimen [see *Clinical Pharmacology (12.3) and Clinical Studies (14)*].

Juvenile Animal Toxicity Data

Oral administration of risdiplam (0, 0.75, 1.5, 2.5 mg/kg/day) to young rats from postnatal day (PND) 4 through PND 31 resulted in decreased growth (body weight, tibia length) and delayed sexual maturation in males at the mid and high dose. The skeletal and body weight deficits persisted after cessation of dosing. Ophthalmic changes consisting of vacuoles in the anterior vitreous were seen at the high dose. Decreases in absolute B lymphocyte counts were observed at all doses after cessation of dosing. Decreases in testis and epididymis weights, which correlated with degeneration of the seminiferous epithelium in the testis, occurred at the mid and high doses; the histopathology findings were reversible, but organ weight persisted after cessation of dosing. Impaired female reproductive performance (decreased mating index, fertility index, and conception rate) was observed at the high dose. A no-effect dose for adverse developmental effects on preweaning rats was not identified. The lowest dose tested (0.75 mg/kg/day) was associated with plasma exposures (AUC) lower than that in humans at the maximum recommended human dose (MRHD) of 5 mg/day.

Oral administration of risdiplam (0, 1, 3, or 7.5 mg/kg/day) to young rats from PND 22 through PND 112 produced a marked increase in micronuclei in the bone marrow, male reproductive organ histopathology (degeneration/necrosis of the seminiferous tubule epithelium,

oligo/aspermia in the epididymis, spermatic granulomas), and adverse effects on sperm parameters (decreased sperm concentration and motility, increased sperm morphology abnormalities) at the highest dose tested. Increases in T lymphocytes (total, helper, and cytotoxic) were observed at the mid and high doses. The reproductive and immune effects persisted after cessation of dosing. The no-effect dose (1 mg/kg/day) for adverse effects on postweaning juvenile rats was associated with plasma exposures (AUC) lower than that in humans at the MRHD.

8.5 Geriatric Use


Clinical studies of EVRYSDI did not include patients aged 65 years and older to determine whether they respond differently from younger adult patients.

11 DESCRIPTION

EVRYSDI for oral solution and EVRYSDI tablets for oral use contain risdiplam, which is a survival of motor neuron 2 (SMN2)-directed RNA splicing modifier.

The chemical name of risdiplam is 7-(4,7-diazaspiro[2.5]octan-7-yl)-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido-4H-[1,2-a]pyrimidin-4-one. Risdiplam has a molecular weight of 401.46 g/mol. Risdiplam demonstrates pH-dependent aqueous solubility; the greatest solubility is at low pH, and solubility decreases with increasing pH. Risdiplam has a pKa₁ of 3.78 (base) and pKa₂ of 6.62 (base).

The molecular formula of risdiplam is C₂₂H₂₃N₇O and the chemical structure is shown below.



EVRYSDI for oral solution is supplied as a powder in an amber glass bottle. Each bottle contains 60 mg of risdiplam. The inactive ingredients of EVRYSDI are: ascorbic acid, disodium edetate dihydrate, isomalt, mannitol, polyethylene glycol 6000, sodium benzoate, strawberry flavor, sucralose, and tartaric acid.

The powder is constituted with purified water to yield 60 mg/80 mL (0.75 mg/mL) of risdiplam after constitution [*see Dosage and Administration (2.4)*].

Each EVRYSDI tablet contains 5 mg of risdiplam. The inactive ingredients of EVRYSDI tablet are colloidal silicon dioxide, crospovidone, mannitol, microcrystalline cellulose, polyethylene glycol 3350, polyvinyl alcohol, sodium stearyl fumarate, strawberry flavor, talc, tartaric acid, titanium dioxide, and yellow iron oxide.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Risdiplam is a survival of motor neuron 2 (SMN2) splicing modifier designed to treat patients with spinal muscular atrophy (SMA) caused by mutations in chromosome 5q that lead to SMN protein deficiency. Using in vitro assays and studies in transgenic animal models of SMA, risdiplam was shown to increase exon 7 inclusion in SMN2 messenger ribonucleic acid (mRNA) transcripts and production of full-length SMN protein in the brain.

In vitro and in vivo data indicate that risdiplam may cause alternative splicing of additional genes, including FOXM1 and MADD. FOXM1 and MADD are thought to be involved in cell

cycle regulation and apoptosis, respectively, and have been identified as possible contributors to adverse effects seen in animals.

12.2 Pharmacodynamics

In clinical trials for infantile-onset SMA and later-onset SMA patients, EVRYSDI led to an increase in SMN protein with a greater than 2-fold median change from baseline within 4 weeks of treatment initiation across all SMA types studied. The increase was sustained throughout the treatment period (of at least 24 months).

Cardiac Electrophysiology

At the maximum recommended dose, clinically significant QTc interval prolongation was not observed.

12.3 Pharmacokinetics

Pharmacokinetics of EVRYSDI have been characterized in healthy adult subjects and in patients with SMA.

After administration of EVRYSDI as an oral solution, pharmacokinetics of risdiplam were approximately linear between 0.6 and 18 mg in a single-ascending-dose study in healthy adult subjects, and between 0.02 and 0.25 mg/kg once daily in a multiple-ascending-dose study in patients with SMA. Following once-daily oral administration of risdiplam in healthy subjects, approximately 3-fold accumulation of peak plasma concentrations (C_{max}) and area under the plasma concentration-time curve (AUC_{0-24h}) was observed. Risdiplam exposures reach steady state 7 to 14 days after once daily administration. EVRYSDI tablet (swallowed whole or dispersed in water) demonstrated comparable bioavailability to EVRYSDI for oral solution in adult healthy volunteers under fasted and fed states.

Absorption

Following oral administration of risdiplam in fasted state, the median time to reach maximum plasma concentration (T_{max}) was 3.26 to 4 hours. The T_{max} was delayed by up to 1 hour in fed state compared to that under fasted state.

Effect of Food

Food (high-fat, high calorie breakfast) had no relevant effect on the exposure of risdiplam. In the clinical efficacy studies (Study 1 and Study 2), risdiplam was administered with a morning meal or after breastfeeding.

Distribution

The apparent volume of distribution at steady state is 190.4 L for a 31.3 kg patient.

Risdiplam is predominantly bound to serum albumin, without any binding to alpha-1 acid glycoprotein, with a free fraction of 11%.

Elimination

The apparent clearance (CL/F) of risdiplam is 2.45 L/h for a 31.3 kg patient. The terminal elimination half-life of risdiplam was approximately 50 hours in healthy adults.

Metabolism

Risdiplam is primarily metabolized by flavin monooxygenase 1 and 3 (FMO1 and FMO3) and also by CYPs 1A1, 2J2, 3A4, and 3A7.

Parent drug was the major component found in plasma, accounting for 83% of drug-related material in circulation. The pharmacologically-inactive metabolite M1 was identified as the major circulating metabolite.

Excretion

Following a dose of 18 mg, approximately 53% of the dose (14% unchanged risdiplam) was excreted in the feces and 28% in urine (8% unchanged risdiplam).

Specific Populations

There were no clinically significant differences in the pharmacokinetics of EVRYSDI based on race or gender. Renal impairment is not expected to alter the exposures to risdiplam.

The impact of geriatric age on the pharmacokinetics of EVRYSDI has not been studied.

Hepatic Impairment

The pharmacokinetics and safety of risdiplam have been studied in subjects with mild or moderate hepatic impairment (as defined by Child-Pugh class A and B, respectively, n = 8 each) compared to subjects with normal hepatic function (n = 10). Following the administration of 5 mg EVRYSDI, the AUC_{inf} and C_{max} of risdiplam were approximately 20% and 5% lower, respectively, in subjects with mild hepatic impairment and were approximately 8% and 20% higher, respectively, in subjects with moderate hepatic impairment, versus matched healthy control subjects. The magnitude of these changes is not considered to be clinically meaningful. The pharmacokinetics and safety in patients with severe hepatic impairment (Child-Pugh class C) have not been studied.

Pediatric Patients

Body weight and age were found to have significant effect on the pharmacokinetics of risdiplam. The estimated exposure (mean AUC_{0-24h}) in pre-symptomatic infants at the age of 1 to 2 months was 2090 ng.h/mL at the recommended dose of 0.15 mg/kg once daily. The estimated exposure for infantile-onset SMA patients (age 2 to 7 months at enrollment) at the recommended dose of 0.2 mg/kg once daily was 1930 ng.h/mL. The estimated exposure for later-onset SMA patients (2 to 25 years old at enrollment) at the recommended dose was 2070 ng.h/mL (0.25 mg/kg once daily for patients with a body weight < 20 kg and 5 mg once daily for patients with a body weight ≥ 20 kg).

No data on risdiplam pharmacokinetics are available in patients less than 16 days of age [*see Use in Specific Populations (8.4)*].

Drug Interaction Studies

Effect of Other Drugs on EVRYSDI

Coadministration of 200 mg itraconazole (a strong CYP3A inhibitor) twice daily with a single 6 mg oral dose of risdiplam did not have a clinically relevant effect on the pharmacokinetics of risdiplam (11% increase in AUC and 9% decrease in C_{max}).

Risdiplam is a weak substrate of human MDR-1 and breast cancer resistant protein (BCRP) transporters in vitro. Human MDR-1 or BCRP inhibitors are not expected to result in a clinically significant increase of risdiplam concentrations.

The effect of omeprazole (an acid-reducing agent/proton pump inhibitor) on risdiplam pharmacokinetics was investigated in healthy subjects. No clinically significant difference was observed in the pharmacokinetics of risdiplam administered as a tablet when used concomitantly with omeprazole. Based on these results, no clinically significant differences are expected with other acid-reducing agents, including H₂-receptor antagonists and antacids.

Effect of EVRYSDI on Other Drugs

Risdiplam and its major circulating metabolite M1 did not induce CYP1A2, 2B6, 2C8, 2C9, 2C19, or 3A4 in vitro. Risdiplam and M1 did not inhibit (reversible or time-dependent

inhibition) any of the CYP enzymes tested (CYP1A2, 2B6, 2C8, 2C9, 2C19, 2D6), with the exception of CYP3A *in vitro*.

EVRYSDI is a weak inhibitor of CYP3A. In healthy adult subjects, administration of EVRYSDI once daily for 2 weeks slightly increased the exposure of midazolam, a sensitive CYP3A substrate (AUC 11%; C_{max} 16%); this increase is not considered clinically relevant. Based on physiologically-based pharmacokinetic (PBPK) modeling, a similar increase is expected in children and infants as young as 2 months of age.

In vitro studies have shown that risdiplam and its major metabolite are not significant inhibitors of human MDR1, organic anion-transporting polypeptide (OATP) 1B1, OATP1B3, organic anion transporter 1 and 3 (OAT 1 and 3) transporters, and human organic cation transporter 2 (OCT2), at clinically relevant concentrations. Risdiplam and its metabolite are, however, *in vitro* inhibitors of the multidrug and toxin extrusion (MATE) 1 and MATE2-K transporters [*see Drug Interactions (7.1)*].

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Risdiplam was not carcinogenic in Tg.rasH2 mice when administered at oral doses of up to 9 mg/kg/day for 26 weeks.

In a 2-year carcinogenicity study in rats, oral administration of risdiplam (0, 0.3, 1, or 3 mg/kg/day) resulted in increased incidences of preputial gland squamous cell carcinomas in males and combined thyroid follicular cell adenomas and carcinomas in females at the highest dose tested. The higher dose not associated with an increase in tumors (1 mg/kg/day) was associated with plasma drug exposures (AUC) similar to that in humans at the maximum recommended human dose (MRHD) of 5 mg/day.

Mutagenesis

Risdiplam was negative in an *in vitro* Ames assay. In an *in vivo* combined bone marrow micronucleus and comet assay in rat, risdiplam was clastogenic, as evidenced by an increase in micronuclei in bone marrow, but was negative in the comet assay. A pronounced increase in bone marrow micronuclei was also observed in toxicity studies in adult and juvenile rats [*see Use in Specific Populations (8.4)*].

Impairment of Fertility

Oral administration of risdiplam to rats for 4 (0, 1, 3, or 9 mg/kg/day) or 26 (0, 1, 3, or 7.5 mg/kg/day) weeks resulted in histopathological effects in the testis (degenerated spermatocytes, degeneration/atrophy of the seminiferous tubules) and epididymis (degeneration/necrosis of ductular epithelium) at the mid and/or high doses. At the high dose in the 26-week study, the testicular lesions persisted to the end of the recovery period, which corresponds, in rat, to approximately one spermatogenic cycle. The no-effect dose for adverse reproductive system effects in adult male rats (1 mg/kg/day) was associated with plasma drug exposures (AUC) similar to that in humans at the maximum recommended human dose (MRHD) of 5 mg/day.

Adverse effects of risdiplam on the testis could not be fully evaluated in the monkey because the majority of monkeys tested were sexually immature. However, oral administration of risdiplam (0, 2, 4, or 6 mg/kg/day) for 2 weeks resulted in histopathological changes in the testis (increases in multinucleate cells, germ cell degeneration) at the highest dose. At the no-effect dose for testicular toxicity in monkeys, plasma exposures were approximately 3 times that in humans at the MRHD.

Oral administration of risdiplam to postweaning juvenile rats resulted in male reproductive toxicity (degeneration/necrosis of the testis seminiferous epithelium with associated oligo/aspermia in the epididymis and abnormal sperm parameters). The no-effect dose for adverse reproductive effects in postweaning male juvenile rats was associated with plasma exposures approximately 4 times that in humans at the MRHD [*see Use in Specific Populations (8.4)*].

13.2 Animal Toxicology and/or Pharmacology

Retinal toxicity

Risdiplam-induced functional and structural retinal abnormalities were seen in animal studies. In a 39-week toxicity study in monkeys, oral administration of risdiplam (0, 1.5, 3, or 7.5/5 mg/kg/day; high dose lowered after 4 weeks) produced functional abnormalities on the electroretinogram (ERG) in all mid- and high-dose animals at the earliest examination time (Week 20). These findings were associated with retinal degeneration, detected by optical coherence tomography (OCT), on Week 22, the first examination time. The retinal degeneration, with peripheral photoreceptor loss, was irreversible. A no-effect dose for the retinal findings (1.5 mg/kg/day) was associated with plasma exposures (AUC) similar to that in humans at the maximum recommended human dose (MRHD) of 5 mg.

Effect on Epithelial Tissues

Oral administration of risdiplam to rats and monkeys resulted in histopathological changes in epithelium of the gastrointestinal (GI) tract (apoptosis/single cell necrosis), lamina propria (vacuolation), the exocrine pancreas (single cell necrosis), the skin, tongue, and larynx (parakeratosis/hyperplasia/degeneration) with associated inflammation. The skin and GI epithelial effects were reversible. The no-effect doses for effects on epithelial tissues in rats and monkeys were associated with plasma exposures (AUC) similar to that in humans at the MRHD.

14 CLINICAL STUDIES

The efficacy of EVRYSDI for the treatment of patients with infantile-onset, later-onset, and pre-symptomatic SMA was evaluated in three clinical studies, Study 1 (NCT02913482) and Study 2 (NCT02908685), and Study 3 (NCT03779334), respectively.

The overall findings of these studies support the effectiveness of EVRYSDI in SMA pediatric and adult patients and appear to support the early initiation of treatment with EVRYSDI.

14.1 Infantile-Onset SMA

Study 1 was an open-label, 2-part study to investigate the efficacy, safety, pharmacokinetics, and pharmacodynamics of EVRYSDI for oral solution in patients with Type 1 SMA (symptom onset between 28 days and 3 months of age). All patients had genetic confirmation of homozygous deletion or compound heterozygosity predictive of loss of function of the SMN1 gene, and two SMN2 gene copies.

Part 1 of Study 1 was designed as a dose-finding study. Part 2 of Study 1 assessed the safety and efficacy of EVRYSDI at 0.20 mg/kg, the recommended dose determined in Part 1 [*see Dosage and Administration (2.4)*]. Patients from Part 1 did not take part in Part 2.

A total of 62 patients with symptomatic Type 1 SMA were enrolled in FIREFISH Part 1 (n = 21) and Part 2 (n = 41), of which 58 patients received the recommended dosage [*see Dosage and Administration (2.1)*]. The median age of onset of clinical signs and symptoms was 1.5 months (range: 0.9 to 3.0 months). The median age at enrollment was 5.6 months (range: 2.2 to 6.9 months), and the median time between onset of symptoms and the first dose was 3.7 months (range 1.0 to 6.0 months). Of these patients, 60% were female, 57% were Caucasian, and 29%

were Asian. The demographics and baseline disease characteristics were comparable between Part 1 and Part 2 of the study.

Effectiveness was established based on the ability to sit without support for at least 5 seconds (as measured by Item 22 of the Bayley Scales of Infant and Toddler Development – Third Edition (BSID-III) gross motor scale) and on the basis of survival without permanent ventilation. Permanent ventilation was defined as requiring a tracheostomy or more than 21 consecutive days of either non-invasive ventilation (≥ 16 hours per day) or intubation, in the absence of an acute reversible event.

The primary endpoint was the proportion of patients with the ability to sit without support for at least 5 seconds (BSID-III gross motor scale, Item 22) after 12 months of treatment in Part 2; 29% of patients (n = 12/41) achieved this milestone.

Other efficacy endpoints of EVRYSDI-treated patients in Study 1 (pooled Part 1 and Part 2) are shown in Table 3.

Table 3 Key Efficacy Results at Month 12 and Month 24 (Study 1, Parts 1 and Part 2)

Efficacy Endpoints	Proportion of Patients Parts 1 & 2 at Month 12	Proportion of Patients Parts 1 & 2 at Month 24
Motor Function and Development Milestones	N = 58 ^a	
BSID-III, Item 22: sitting without support for at least 5 seconds	32.8%	60.3%
Survival and Event-Free Survival	N = 62 ^b	
Alive without Permanent Ventilation	87.1%	83.8%

^a Results were pooled from all patients who received the recommended dose of risdiplam (all patients in Part 2 and those in the high-dose cohort of Part 1; n = 58).

^b Results were pooled from all patients who received any dose of risdiplam in Part 1 and Part 2 (n = 62).

At Month 24, 40% (23/58) of patients who received the recommended dose achieved sitting without support for 30 seconds (BSID-III, Item 26). In addition at Month 24, patients continued to achieve additional motor milestones; 28% (16/58) of patients achieved a standing measure (16% [9/58] supporting weight and 12% [7/58] standing with support), as measured by Section 2 of the Hammersmith Infant Neurological Examination (HINE-2) which assesses motor milestones.

The proportion of patients alive without permanent ventilation (event-free survival) was 84% for all patients at Month 24 (Table 3). Out of 62 patients, 6 infants died (4 within the first 3 months following study enrollment) and one additional patient withdrew from treatment and died 3.5 months later. Four patients required permanent ventilation by Month 24. These results indicate a clinically meaningful deviation from the natural history of untreated infantile-onset SMA. As described in the natural history of untreated infantile-onset SMA, patients would not be expected to attain the ability to sit independently, and no more than 25% of these patients would be expected to survive without permanent ventilation beyond 14 months of age.

14.2 Later-Onset SMA

Study 2 was a 2-part, multicenter trial to investigate the efficacy, safety, pharmacokinetics, and pharmacodynamics of EVRYSDI for oral solution in patients diagnosed with SMA Type 2 or Type 3. Part 1 of Study 2 was dose-finding and exploratory in 51 patients (14% ambulatory). Part 2 was randomized, double-blind, placebo-controlled, and is described below.

The primary endpoint in Study 2 Part 2 was the change from baseline to Month 12 in the Motor Function Measure 32 (MFM32) score. A key secondary endpoint was the proportion of patients with a 3-point or greater change from baseline to Month 12 in the MFM32 total score. The MFM32 measures motor function abilities that relate to daily functions. The total MFM32 score is expressed as a percentage (range: 0 to 100) of the maximum possible score, with higher scores indicating greater motor function. Another key secondary endpoint was the Revised Upper Limb Module (RULM). The RULM is a tool used to assess motor performance of the upper limb in SMA patients. It tests proximal and distal motor functions of the arm. The total score ranges from 0 (all the items cannot be performed) to 37 (all the activities are achieved fully without any compensatory maneuvers).

Study 2 Part 2 enrolled 180 non-ambulatory patients with Type 2 (71%) or Type 3 (29%) SMA. Patients were randomized 2:1 to receive EVRYSDI at the recommended dosage [*see Dosage and Administration (2.1)*] or placebo. Randomization was stratified by age group (2 to 5, 6 to 11, 12 to 17, or 18 to 25 years of age).

The median age of patients at the start of treatment was 9.0 years (range: 2 to 25), and the median time between onset of initial SMA symptoms and first treatment was 102.6 months (range: 1 to 275). Of the 180 patients included in the trial, 51% were female, 67% were Caucasian, and 19% were Asian. At baseline, 67% of patients had scoliosis (32% of them with severe scoliosis). Patients had a mean baseline MFM32 score of 46.1, and RULM score of 20.1. Overall baseline demographic characteristics were reasonably balanced between the treatment groups (EVRYSDI and placebo), with the exception of scoliosis (63% in the EVRYSDI arm vs. 73% in the placebo group).


The primary analysis on the change from baseline in MFM32 total score at Month 12 showed a clinically meaningful and statistically significant difference between patients treated with EVRYSDI and placebo. The results of the primary analysis and key secondary endpoints are shown in Table 4 and Figure 1.

Table 4 Summary of Efficacy in Patients with Later-Onset SMA at Month 12 of Treatment (Study 2 Part 2)

Endpoint	EVRYSDI (N = 120)	Placebo (N = 60)
Primary Endpoint:		
Change from baseline in total MFM32 score at Month 12, LS means (95% CI) ^{1,2,3}	1.36 (0.61, 2.11)	-0.19 (-1.22, 0.84)
Difference from Placebo, Estimate (95% CI) ¹ p-value	1.55 (0.30, 2.81) 0.0156	
Secondary Endpoints:		
Proportion of patients with a change from baseline MFM32 total score of 3 or more at Month 12 (95% CI) ^{2,3}	38.3% (28.9, 47.6)	23.7% (12.0, 35.4)
Odds ratio for overall response (95% CI) adjusted ⁴ (unadjusted) p-value ⁵	2.35 (1.01, 5.44) 0.0469 (0.0469)	
Change from baseline in total score of RULM at Month 12, LS means (95% CI) ^{1, 6}	1.61 (1.00, 2.22)	0.02 (-0.83, 0.87)
Difference from Placebo, Estimate (95% CI) adjusted ⁴ (unadjusted) p-value ¹	1.59 (0.55, 2.62) 0.0469 (0.0028)	

- ¹. The Mixed Model Repeated Measure (MMRM) analysis included the change from baseline total score as the dependent variable and as independent variables the baseline total score, treatment group, time, treatment-by-time interaction, and the randomization stratification variable of age group (2 to 5, 6 to 11, 12 to 17, 18 to 25).
- ². The MFM total score was calculated according to the user manual, expressed as a percentage of the maximum score possible for the scale (i.e., sum of the 32 item scores divided by 96 and multiplied by 100).
- ³. Based on the missing data rule for MFM32, 6 patients were excluded from the analysis (EVRYSDI n = 115; placebo control n = 59).
- ⁴. The adjusted p-value was obtained for the endpoints included in the hierarchical testing and was derived based on all the p-values from endpoints in order of the hierarchy up to the current endpoint.
- ⁵. The logistic regression analysis included the baseline total score, treatment and age group as independent variables.
- ⁶. Based on the missing data rule for RULM, 3 patients were excluded from the analysis (EVRYSDI n = 119; placebo control n = 58).

Figure 1 Mean Change from Baseline in Total MFM32 Score Over 12 Months (Study 2 Part 2)^{1,2}



¹ Error bars denote the 95% confidence interval.

² The MFM total score was calculated according to the user manual, expressed as a percentage of the maximum score possible for the scale (i.e., sum of the 32 item scores divided by 96 and multiplied by 100).

14.3 Pre-Symptomatic SMA

Study 3 was an open-label, single-arm, multicenter clinical study to investigate the efficacy, safety, pharmacokinetics, and pharmacodynamics of EVRYSDI in infants up to 6 weeks of age (at first dose) who have been genetically diagnosed with SMA but do not yet present with symptoms.

The efficacy in pre-symptomatic SMA patients was evaluated at Month 12 in 26 patients treated with EVRYSDI in Study 3: 8 patients had 2 copies of the *SMN2* gene, 13 patients had 3 copies, and 5 patients had 4 or more copies. The median age of these patients at first dose was 25 days (range: 16 to 41), 62% were female, and 85% were Caucasian. The primary efficacy population (N = 5) included patients with 2 *SMN2* copies and a baseline CMAP amplitude ≥ 1.5 mV.

The primary efficacy endpoint was the proportion of patients with the ability to sit without support for at least 5 seconds (BSID-III gross motor scale, Item 22) at Month 12. This milestone was achieved by 80% (4/5) of patients in the primary efficacy population. This milestone was also achieved by 87.5% (7/8) of all patients with 2 copies of *SMN2* and 96.2% (25/26) of patients in the full treated population.

At Month 12, 80.8% (21/26) of patients in the full treated population achieved sitting without support for 30 seconds (BSID-III, Item 26). Of the 26 patients treated with EVRYSDI, 25 patients had motor milestones measured by the HINE-2 at Month 12. Of these, 24 (96%) achieved sitting (23 patients could pivot/rotate and 1 achieved stable sit); 21 (84%) could stand (13 patients could stand unaided and 8 could stand with support); and 12 (48%) could walk independently. Seven patients were not tested for walking at Month 12. All 26 patients were alive at 12 months without permanent ventilation.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 EVRYSDI for Oral Solution

How Supplied

Each amber glass bottle of EVRYSDI for oral solution is packaged with a bottle adapter, two 1 mL reusable oral syringes, two 6 mL reusable oral syringes, and one 12 mL reusable oral syringe. EVRYSDI for oral solution is a light yellow, pale yellow, yellow, greyish yellow, greenish yellow, or light green powder. Each bottle contains 60 mg of risdiplam (NDC 50242-175-07).

Storage and Handling

Store the dry powder at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C to 30°C (59°F to 86°F) [*see USP controlled room temperature*]. Keep in the original carton.

Keep the constituted oral solution of EVRYSDI in the original amber bottle to protect from light. Store in a refrigerator at 2°C to 8°C (36°F to 46°F) [*see Dosage and Administration (2.4)*].

16.2 EVRYSDI Tablets

How Supplied

Pale yellow film-coated tablet, round and curved, with EVR debossed on one side; available in HDPE bottles of 30 tablets with a child-resistant cap (NDC 50242-202-01).

Storage and Handling

Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C to 30°C (59°F to 86°F) [*see USP Controlled Room Temperature*]. Keep in the original bottle. Keep the bottle tightly closed in order to protect from moisture.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information and Instructions for Use).

Pregnancy and Fetal Risk

Inform pregnant women and women of reproductive potential that, based on animal studies, EVRYSDI may cause fetal harm [*see Use in Specific Populations (8.1)*].

Discuss with women of childbearing age whether they are pregnant, might be pregnant, or are trying to become pregnant.

Advise women of childbearing potential to use effective contraception during treatment with EVRYSDI and for at least 1 month after stopping EVRYSDI.

Advise a female patient to immediately inform the prescriber if she is pregnant or planning to become pregnant [*see Use in Specific Populations (8.3)*].

Pregnancy Registry

Encourage patients to enroll in the EVRYSDI Pregnancy Registry if they become pregnant while taking EVRYSDI [*see Use in Specific Populations (8.1)*].

Potential Effects on Male Fertility

Advise male patients that their fertility may be compromised while on treatment with EVRYSDI [*see Use in Specific Populations (8.3)*].

Instructions for Preparation of Oral Solution

Advise patients/caregivers to ensure that EVRYSDI oral solution is in liquid form when received from the pharmacy.

Instruct patients/caregivers to take EVRYSDI oral solution with or without food or before or after breastfeeding at approximately the same time each day. However, instruct caregivers not to mix EVRYSDI with formula or milk.

Instruct patients/caregivers to take EVRYSDI oral solution immediately after it is drawn up into the reusable oral syringe [see *Dosage and Administration (2.2)*].

Instructions for EVRYSDI Tablets

Advise patients/caregivers to swallow EVRYSDI tablets whole with water. Do not chew, cut, or crush the tablets.

Alternatively, the tablet can be dispersed in one teaspoon (5 mL) of room temperature non-chlorinated drinking water (e.g., filtered water) and taken immediately. EVRYSDI tablets must not be dispersed in any liquid other than non-chlorinated drinking water. Instruct the patient/caregivers that the dispersion must be administered within 10 minutes of adding non-chlorinated drinking water, or it must be discarded [see *Dosage and Administration (2.2)*].

Advise patients/caregivers to wash their hands before and after preparing or taking EVRYSDI tablets.

Advise patients/caregivers to avoid getting the dispersed tablet on their skin or in their eyes. Advise patients/caregivers to wash the area with soap and water if the dispersed tablet gets on the skin. Advise patients/caregivers to rinse their eyes with water if the dispersed tablet gets in the eyes.

Advise patients/caregivers to use a dry paper towel to dry the area if the dispersion is spilled and then clean with soap and water. Advise patients/caregivers to throw the paper towel away in the trash and wash their hands with soap and water.

EVRYSDI® [risdiplam]

Distributed by:
Genentech USA, Inc.
A Member of the Roche Group
1 DNA Way
South San Francisco, CA 94080-4990

EVRYSDI is a registered trademark of Genentech, Inc.
©2026 Genentech, Inc. All rights reserved.

Patient Information

**EVERYSDI® [ev-RIZ-dee]
(risdiplam)
for oral solution**

**EVERYSDI® [ev-RIZ-dee]
(risdiplam)
tablets, for oral use**

What is EVERYSDI?

- EVERYSDI is a prescription medicine used to treat spinal muscular atrophy (SMA) in children and adults.

Before taking EVERYSDI, tell your healthcare provider about all of your medical conditions, including if you:

- are pregnant or plan to become pregnant. If you are pregnant, or are planning to become pregnant, ask your healthcare provider for advice before taking this medicine. EVERYSDI may harm your unborn baby.
- are a woman who can become pregnant:
 - Before you start your treatment with EVERYSDI, your healthcare provider may test you for pregnancy. Because EVERYSDI may harm your unborn baby, you and your healthcare provider will decide if taking EVERYSDI is right for you during this time.
 - Talk to your healthcare provider about birth control methods that may be right for you. Use birth control while on treatment and for at least 1 month after stopping EVERYSDI.
 - **Pregnancy Registry.** There is a pregnancy registry for women who take EVERYSDI during pregnancy. The purpose of this registry is to collect information about the health of the pregnant woman and her baby. If you are pregnant or become pregnant while receiving EVERYSDI, tell your healthcare provider right away. Talk to your healthcare provider about registering with the EVERYSDI Pregnancy Registry. Your healthcare provider can enroll you in this registry or you can enroll by calling 1-833-760-1098 or visiting <https://www.evrysdipregnancyregistry.com>.
- are an adult male planning to have children: EVERYSDI may affect a man's ability to have children (fertility). If this is of concern to you, make sure to ask a healthcare provider for advice.
- are breastfeeding or plan to breastfeed. It is not known if EVERYSDI passes into breast milk and may harm your baby. If you plan to breastfeed, discuss with your healthcare provider about the best way to feed your baby while on treatment with EVERYSDI.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Keep a list of them to show your healthcare provider, including your pharmacist, when you get a new medicine.

How should I take EVERYSDI?

- For infants and children, your healthcare provider will determine the daily dose of EVERYSDI needed based on your child's age and weight. For adults, take 5 mg of EVERYSDI daily.
- Your healthcare provider will either prescribe:
EVERYSDI for oral solution
Or
EVERYSDI tablet
- Your healthcare provider will tell you how long you or your child needs to take EVERYSDI. Do not stop treatment with EVERYSDI unless your healthcare provider tells you to.
 - Take EVERYSDI exactly as your healthcare provider tells you to take it. Do not change the dose without talking to your healthcare provider.
- Avoid getting EVERYSDI on your skin or in your eyes. If it gets on your skin, wash the area with soap and water. If it gets in your eyes, rinse your eyes with water.

If you are taking EVERYSDI for oral solution:

- **See the detailed Instructions for Use that comes with it for information on how to take or give EVERYSDI for oral solution.**
- You should receive EVERYSDI from the pharmacy as a liquid that can be given by mouth or through a feeding tube. The liquid solution is prepared by your pharmacist or other healthcare provider. If the medicine in the bottle is a powder, **do not use it**. Contact your pharmacist for a replacement.
- Take EVERYSDI one time daily with or without a meal at about the same time each day. Drink water afterwards to make sure EVERYSDI has been completely swallowed.

- In infants who are breastfed, EVRYSDI can be given before or after breastfeeding.
- **Do not** mix EVRYSDI with formula or milk.
- If you are unable to swallow and have a nasogastric (NG-tube) or gastrostomy tube (G-tube), EVRYSDI for oral solution can be given through the tube.

Reusable Oral Syringes for EVRYSDI for Oral Solution

- Your pharmacist will provide you with the reusable oral syringes that are needed for taking your medicine and explain how to use them. You should receive 1 or 2 identical oral syringes depending on your prescribed daily dose.
- From the bottle, draw up (measure) the dose of EVRYSDI with these syringes, as they are made to protect the medicine from light.
- Take EVRYSDI right away after it has been drawn into the syringe. **Do not** store the EVRYSDI solution in the syringe. If EVRYSDI is not taken within 5 minutes of when it is drawn up, throw away the solution by pressing the plunger and prepare a new dose with the same syringe.
- **Do not** throw the syringes away because they are reusable.
- Wash the syringes per instructions after use.
- Contact your healthcare provider or pharmacist if your oral syringes are lost or damaged.

If you are taking EVRYSDI Tablets:

- **See the detailed Instructions for Use that comes with it for information on how to take or give EVRYSDI tablets.**
- **Wash your hands before and after preparing or taking EVRYSDI tablets.**
- Take EVRYSDI one time daily with or without a meal at about the same time each day.
- **Do not** chew, cut, or crush the tablet.
- Swallow the EVRYSDI tablet whole with some water.
- If you cannot swallow tablets whole or if you have a nasogastric (NG-tube) or gastrostomy (G-tube), see the detailed Instructions for Use that comes with it for information on how to take or give EVRYSDI tablets.

If you miss a dose of EVRYSDI:

- If you remember the missed dose within 6 hours of when you normally take EVRYSDI, then take the dose. Continue taking EVRYSDI at your usual time the next day.
- If you remember the missed dose more than 6 hours after you normally take EVRYSDI, skip the missed dose. Take your next dose at your usual time the next day.
- If you do not fully swallow the dose, or you vomit after taking a dose, **do not take** another dose of EVRYSDI to make up for that dose. Wait until the next day to take the next dose at your usual time.

What are the possible side effects of EVRYSDI?

The most common side effects of EVRYSDI include:

- **For later-onset SMA:**
 - fever
 - diarrhea
 - rash
- **For infantile-onset SMA:**
 - fever
 - runny nose, sneezing, and sore throat (upper respiratory infection)
 - constipation
 - diarrhea
 - lung infection (lower respiratory infection)
 - vomiting
 - rash
 - cough

These are not all of the possible side effects of EVRYSDI. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store EVRYSDI?

EVRYSDI for Oral Solution:

- Store EVRYSDI in the refrigerator between 36°F to 46°F (2°C to 8°C). Do not freeze.

- If necessary, EVRYSDI can be kept at room temperature up to 104°F (up to 40°C) for a combined total of 5 days. EVRYSDI can be removed from, and returned to, a refrigerator. The total combined time out of refrigeration should not be more than 5 days.
- Keep EVRYSDI in an upright position in the original amber bottle to protect from light.
- Throw away (discard) any unused portion of EVRYSDI 64 days after it is mixed by the pharmacist (constitution) or if EVRYSDI has been kept at room temperature (below 104°F [40°C]) for more than a total combined time of 5 days. Discard EVRYSDI if it has been kept above 104°F (40°C). Please see the Discard After date written on the bottle label. (See the **Instructions for Use** that comes with EVRYSDI for oral solution).

EVRYSDI Tablets:

- Store at room temperature between 68°F to 77°F (20°C to 25°C).
- Keep the tablets in their original bottle with the cap tightly closed to protect from moisture.

Keep EVRYSDI, all medicines and syringes out of the reach of children.

General information about the safe and effective use of EVRYSDI.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use EVRYSDI for a condition for which it was not prescribed. Do not give EVRYSDI to other people, even if they have the same symptoms you have. It may harm them. You can ask your pharmacist or healthcare provider for information about EVRYSDI that is written for health professionals.

What are the ingredients in EVRYSDI?

Active ingredient: risdiplam

Inactive ingredients:

EVRYSDI for Oral Solution: ascorbic acid, disodium edetate dihydrate, isomalt, mannitol, polyethylene glycol 6000, sodium benzoate, strawberry flavor, sucralose, and tartaric acid.

EVRYSDI Tablets: colloidal silicon dioxide, crospovidone, mannitol, microcrystalline cellulose, polyethylene glycol 3350, polyvinyl alcohol, sodium stearyl fumarate, strawberry flavor, talc, tartaric acid, titanium dioxide, and yellow iron oxide.

Distributed by: **Genentech USA, Inc.**, A Member of the Roche Group, 1 DNA Way, South San Francisco, CA 94080-4990

EVRYSDI is a registered trademark of Genentech, Inc.

©2026 Genentech, Inc. All rights reserved.

For more information, go to www.EVRYSDI.com or call 1-833-387-9734.