



Welcome



A guide to getting started with Evrysdi tablets

Liz,
living with
Type 3 SMA

Joyful,
daughter
to Liz

Patients featured in this guide take Evrysdi, but may not be taking the tablet formulation.

What is Evrysdi?

Evrysdi is a prescription medicine used to treat spinal muscular atrophy (SMA) in children and adults.

Important Safety Information

- **Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:**
 - are pregnant or plan to become pregnant, as Evrysdi may harm your unborn baby. Ask your healthcare provider for advice before taking this medicine

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).



Welcome to the Evrysdi community!

Where To Start

Obtaining Evrysdi

Storing Evrysdi

Taking Evrysdi

Support

FAQs

Important Safety Information

Antoine, partner to Janelle

Adriane, mom to Shaniqua

Janelle, living with Type 2 SMA

Shaniqua, living with Type 3 SMA


This guide is here to help you get started with treatment. On the pages that follow, you'll find answers to your questions and helpful resources.

Important Safety Information (continued)

- **Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:**
 - are a woman who can become pregnant:
 - Before you start your treatment with Evrysdi, your healthcare provider may test you for pregnancy
 - Talk to your healthcare provider about birth control methods that may be right for you. Use birth control while on treatment and for at least 1 month after stopping Evrysdi

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).





Alyson,
Shaniqua's
PAL

Shaniqua,
living with
Type 3 SMA

Where to start: your local contact

The PAL (Partnership and Access Liaison)* team are local points of contact at Genentech, Inc., to answer your questions and offer support and guidance.

A PAL can:

- ✓ **Provide** in-person or virtual support
- ✓ **Answer** questions about Evrysdi
- ✓ **Explain** insurance coverage and potential financial support options
- ✓ **Refer** you to helpful resources

Don't hesitate to reach out!

Get connected at [Evrysdi.com/PAL](https://www.evrysdi.com/PAL) (833) 387-9734
(Monday-Friday, 9am to 8 pm ET)

*MySMA Support™, including the PAL, does not provide medical advice and is not a substitute for your medical team. Your healthcare provider should always be your main resource for any questions about your health and medical care.

Important Safety Information (continued)

- **Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:**
 - are a woman who can become pregnant:
 - **Pregnancy Registry.** There is a pregnancy registry for women who take Evrysdi during pregnancy. The purpose of this registry is to collect information about the health of the pregnant woman and her baby. If you are pregnant or become pregnant while receiving Evrysdi, tell your healthcare provider right away. Talk to your healthcare provider about registering with the Evrysdi Pregnancy Registry. Your healthcare provider can enroll you in this registry or you can enroll by calling [1-833-760-1098](tel:1-833-760-1098) or visiting <https://www.evrysdi.com/pregnancyregistry.com>.

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).






Where to start: a few things to note



- ✓ Evrysdi is delivered directly to your door
- ✓ A specialty pharmacy* will call you to schedule a time for your delivery



- ✓ Each delivery includes 1 bottle of Evrysdi tablets that will last you 30 days



- ✓ Evrysdi tablets can be stored at room temperature (68°F to 77°F, 20°C to 25°C), so you have flexibility in how and where you take them

*Specialty pharmacies are not part of Genentech and maintain independence in their operations and in their role as a healthcare provider.

Important Safety Information (continued)

- **Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:**
 - are an adult male. Evrysdi may affect a man's ability to have children (fertility). Ask a healthcare provider for advice before taking this medicine
 - are breastfeeding or plan to breastfeed. It is not known if Evrysdi passes into breast milk and may harm your baby


Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).





Jim,
living with
Type 3 SMA

Where to start: a few things to note (continued)



- ✓ You can swallow Evrysdi tablets whole with water or mix them with room temperature non-chlorinated drinking water (eg, filtered water, bottled water), depending on which is easier for you
- ✓ Keep room temperature bottled water on hand if you want to prepare Evrysdi as a mixture (don't mix it with any other liquid than bottled water)*
- ✓ Never prepare the Evrysdi tablet mixture in sunlight



- ✓ It can be helpful to set daily dose reminders on your phone or calendar
- ✓ You can also ask your PAL for a Dose Tracking Calendar and refill tips



- ✓ Financial support may be available to you
- ✓ You can find more information at [Evrysdi.com/assistance](https://www.evrysdi.com/assistance)

These are just a few elements to keep in mind. For more details, please read the Dosage and Administration section of the Prescribing Information that came with your prescription.

*Evrysdi must be taken within 10 minutes of being mixed with bottled water.

Important Safety Information (continued)

- **Tell your healthcare provider about all the medicines you take**
- If you were prescribed Evrysdi for Oral Solution, you should receive Evrysdi from the pharmacy as a liquid. If the medicine in the bottle is a powder, **do not use it**. Contact your pharmacist for a replacement
- Avoid getting Evrysdi on your skin or in your eyes. If Evrysdi gets on your skin, wash the area with soap and water. If Evrysdi gets in your eyes, rinse your eyes with water

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).

Where To Start

Obtaining Evrysdi

Storing Evrysdi

Taking Evrysdi

Support

FAQs

Important Safety Information

Jenn,
Janelle's
PAL

Janelle,
living with
Type 2 SMA

Where to start: when you have questions

FOR QUESTIONS ABOUT...

GO TO...

What to expect when starting treatment, your dose, potential side effects, or your results with Evrysdi

✓ **Your healthcare provider**

Your delivery of Evrysdi

✓ **The specialty pharmacy**
(the company that will call you to schedule your delivery)

Potential financial support options, storing or taking your medicine, planning for travel, helpful resources

✓ **MySMA Support* or a PAL**
(833) 387-9734, Monday-Friday 9 AM to 8 PM ET.
You can also ask your healthcare provider or specialty pharmacy for help with these matters

*Enrollment in MySMA Support through the Evrysdi Start Form is mandatory to receive assistance through the program. Participation in MySMA Support is not necessary to receive treatment with Evrysdi.

Important Safety Information (continued)

• **The most common side effects of Evrysdi include:**

- For later-onset SMA: fever, diarrhea, rash
- For infantile-onset SMA: fever; diarrhea; rash; runny nose, sneezing and sore throat (upper respiratory infection); lung infection (lower respiratory infection); constipation; vomiting; cough

These are not all of the possible side effects of Evrysdi. For more information on the risk and benefits profile of Evrysdi, ask your healthcare provider or pharmacist.

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).





Shaniqua,
living with
Type 3 SMA

**Adriane, mom
to Shaniqua**

“ I love to hang out with my friends, cook, catch a movie, or go out to eat. Evrysdi just works in my life because **I’m not confined by taking my medication in one place.** ”

-Shaniqua, living with Type 3 SMA

Obtaining Evrysdi: your delivery

Evrysdi is delivered directly to your door by a specialty pharmacy.* Evrysdi is not available at retail pharmacies.

- You will receive a call from the specialty pharmacy to welcome you and schedule a delivery time that works best for you; your preferred delivery time will be confirmed each month
- Each delivery includes a 30-day supply of medication



To help track your refills, ask your PAL for a Dose Tracking Calendar and refill tips.

If you have **questions about your delivery**, call your specialty pharmacy.

*Specialty pharmacies are not part of Genentech and maintain independence in their operations and in their role as a healthcare provider.

Important Safety Information (continued)

- **Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:**
 - are pregnant or plan to become pregnant, as Evrysdi may harm your unborn baby. Ask your healthcare provider for advice before taking this medicine
 - are a woman who can become pregnant:
 - Before you start your treatment with Evrysdi, your healthcare provider may test you for pregnancy
 - Talk to your healthcare provider about birth control methods that may be right for you. Use birth control while on treatment and for at least 1 month after stopping Evrysdi

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).



Obtaining Evrysdi: your delivery (continued)

Where To
Start

Obtaining
Evrysdi

Storing
Evrysdi

Taking
Evrysdi


Support

FAQs

Important
Safety
Information

Each package includes:

The package you receive includes
a bottle that holds a 30-day
supply of Evrysdi tablets



Discard After
date



Your shipment will also include tips for taking and giving Evrysdi tablets in the
Dosage and Administration section of the Prescribing Information.

Important Safety Information (continued)

- **Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:**
 - are a woman who can become pregnant:
 - **Pregnancy Registry.** There is a pregnancy registry for women who take Evrysdi during pregnancy. The purpose of this registry is to collect information about the health of the pregnant woman and her baby. If you are pregnant or become pregnant while receiving Evrysdi, tell your healthcare provider right away. Talk to your healthcare provider about registering with the Evrysdi Pregnancy Registry. Your healthcare provider can enroll you in this registry or you can enroll by calling **1-833-760-1098** or visiting **<https://www.evrysdipregnancyregistry.com>**.

Please see additional **Important Safety Information** throughout and full **Prescribing Information**.



Storing Evrysdi: flexibility that fits your lifestyle

Where To Start

Obtaining Evrysdi

Storing Evrysdi

Taking Evrysdi

Support

FAQs

Important Safety Information

No need to plan your day around treatment. Whether at work or on vacation, you can travel with Evrysdi tablets without the need for a cooler bag. See below for additional storage information.

Storing your medicine

- Evrysdi tablets should be kept at room temperature (68°F to 77°F, 20°C to 25°C)
- Do not take or give Evrysdi if it has been exposed to temperatures above 86°F (30°C)
- The tablets are sensitive to moisture, so don't transfer them to a pill organizer. Keep them in the bottle they came in and remember to close it tightly
- Keep Evrysdi and all medicines out of the sight and reach of children

Jose, living with Type 3 SMA



Important Safety Information (continued)

- **Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:**
 - are an adult male. Evrysdi may affect a man's ability to have children (fertility). Ask a healthcare provider for advice before taking this medicine
 - are breastfeeding or plan to breastfeed. It is not known if Evrysdi passes into breast milk and may harm your baby

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).





Taking Evrysdi: more freedom in how and where you take it

Each Evrysdi tablet already contains your dose. You can take it with or without food, so you don't need to plan around meals. Just remember to take it at the same time every day.

Be sure to tell your healthcare provider if you or the person in your care is taking:

- any other medications, vitamins, or dietary supplements, including any stool softeners
- additional fiber that may impact bowel movements, as these may need to be adjusted prior to starting Evrysdi



Something to know:

Evrysdi tablets deliver the same demonstrated results and safety as the liquid form



Where To Start

Obtaining Evrysdi

Storing Evrysdi

Taking Evrysdi

Support

FAQs

Important Safety Information


Preparing to take your tablet

Before you take your first dose, you'll want to make sure you have been shown how to do so, and know where to go if you need a refresher.

- Your healthcare provider, pharmacist, or nurse will show you how to prepare and take Evrysdi tablets. Do not do so until your healthcare team has walked you through the steps
- Please read the Dosage and Administration section of the Prescribing Information that comes with your prescription before taking Evrysdi tablets for the first time
- **Watch a video** on how to take Evrysdi tablets



Wash your hands, then push and twist the child resistant cap to open.



Take 1 tablet out of the bottle and close tightly to protect the remaining medication from moisture. Wash your hands with soap and water once you're done.

Important Safety Information (continued)

- **Tell your healthcare provider about all the medicines you take**
- If you were prescribed Evrysdi for Oral Solution, you should receive Evrysdi from the pharmacy as a liquid. If the medicine in the bottle is a powder, **do not use it**. Contact your pharmacist for a replacement
- Avoid getting Evrysdi on your skin or in your eyes. If Evrysdi gets on your skin, wash the area with soap and water. If Evrysdi gets in your eyes, rinse your eyes with water

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).






Taking Evrysdi tablets

If you or the person in your care are planning to swallow an Evrysdi tablet whole, follow this step:



Swallow the tablet whole with water.
Do not chew, cut, or crush the tablet.
Do not swallow with any liquids other than water.

If you're planning on mixing the tablet with bottled water, follow these steps:



Pour 1 teaspoon (5 mL) of room temperature bottled water into a cup.
Add 1 tablet to the water.



Gently swirl the mixture in the cup around until the tablet is fully mixed.
This can take up to 3 minutes. It is normal for the mixture to look a little cloudy.



Drink the mixture within 10 minutes of adding bottled water to the tablet.
If more than 10 minutes have passed, throw the mixture away according to your local requirements and make a new dose.



Refill the cup with at least 1 tablespoon (15 mL) of bottled water and swirl it around to get any extra medicine left in the cup.



Drink the last tablespoon of mixture immediately.

If you spill the Evrysdi tablet mixture, dry the area with a dry paper towel, then clean it with soap and water. Throw the paper towel away and wash your hands with soap and water

Important Safety Information (continued)

•The most common side effects of Evrysdi include:

- For later-onset SMA: fever, diarrhea, rash
- For infantile-onset SMA: fever; diarrhea; rash; runny nose, sneezing and sore throat (upper respiratory infection); lung infection (lower respiratory infection); constipation; vomiting; cough

These are not all of the possible side effects of Evrysdi. For more information on the risk and benefits profile of Evrysdi, ask your healthcare provider or pharmacist.

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).



Where To
Start

Obtaining
Evrysdi

Storing
Evrysdi

Taking
Evrysdi

Support

FAQs

Important
Safety
Information



Where To
Start

Obtaining
Evrysdi

Storing
Evrysdi

Taking
Evrysdi

Support

FAQs

Important
Safety
Information



“ I think choice is always important in any situation, but most certainly when it comes to your body. Evrysdi for me is taking action.

-Angela, living with
Type 2 SMA

Taking Evrysdi: important reminders

WHAT TO DO:

If the medication is vomited, spit up, or not fully swallowed

- Do not take or give any more medication that day
- Wait until the following day to take or give the next dose at the usual dosing time

If a dose is missed

- The missed dose should be taken or given as soon as possible, but no more than 6 hours after the usual dosing time
- If more than 6 hours have passed, skip the missed dose altogether and take or give the next dose the following day

Click [here](#) to find a video demonstrating how to take Evrysdi

Important Safety Information (continued)

- **Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:**
 - are pregnant or plan to become pregnant, as Evrysdi may harm your unborn baby. Ask your healthcare provider for advice before taking this medicine
 - are a woman who can become pregnant:
 - Before you start your treatment with Evrysdi, your healthcare provider may test you for pregnancy
 - Talk to your healthcare provider about birth control methods that may be right for you. Use birth control while on treatment and for at least 1 month after stopping Evrysdi

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).



Taking Evrysdi: cleaning up

How to dispose of Evrysdi

- Follow your local guidelines for proper disposal of medication. Visit www.fda.gov/drugs/ensuring-safe-use-medicine/safe-disposal-medicines or contact your healthcare provider for more information



If the Evrysdi tablet mixture gets on your skin, wash the area with soap and water. If the Evrysdi tablet mixture gets in your eyes, rinse your eyes with water.

If you spill the Evrysdi tablet mixture, dry the area with a paper towel, then clean it with soap and water. Throw the paper towel away and wash your hands with soap and water.

If there is something wrong with your delivery of Evrysdi tablets

- If you receive an incomplete order, a broken bottle, or medication in powder or liquid form, do not take the medication you received, and contact your specialty pharmacy immediately
- Contact your specialty pharmacy with questions about shipping and packaging

Store the bottle out of sight and reach of children. Call Genentech Product Complaints at [1-800-334-0290](tel:1-800-334-0290) to coordinate your free replacement and to receive further instructions.

Important Safety Information (continued)

- **Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:**
 - are a woman who can become pregnant:
 - **Pregnancy Registry.** There is a pregnancy registry for women who take Evrysdi during pregnancy. The purpose of this registry is to collect information about the health of the pregnant woman and her baby. If you are pregnant or become pregnant while receiving Evrysdi, tell your healthcare provider right away. Talk to your healthcare provider about registering with the Evrysdi Pregnancy Registry. Your healthcare provider can enroll you in this registry or you can enroll by calling [1-833-760-1098](tel:1-833-760-1098) or visiting <https://www.evrysdipregnancyregistry.com>.

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).



Taking Evrysdi: dos and don'ts

DO

- ✓ Do take the dose at the same time every day
- ✓ Do keep the bottle tightly closed in order to protect the Evrysdi tablets from moisture
- ✓ Do store Evrysdi tablets at room temperature (68°F to 77°F, 20°C to 25°C)
- ✓ Do check that the tablets are not expired before use so you don't use expired medication
- ✓ Do ask your specialty pharmacy for the Discard After date if it is not written on the bottle label
- ✓ Do swallow the Evrysdi tablet whole with water (and not other liquids)
- ✓ Do mix 1 Evrysdi tablet with at least 1 teaspoon (5 mL) of room temperature bottled water when preparing the mixture



If you have questions about how to take Evrysdi, contact your healthcare provider. A PAL may also be able to answer questions about Evrysdi; to connect with a PAL, call **(833) 387-9734**, Monday-Friday, 9 AM to 8 PM ET

Important Safety Information (continued)

- **Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:**
 - are an adult male. Evrysdi may affect a man's ability to have children (fertility). Ask a healthcare provider for advice before taking this medicine
 - are breastfeeding or plan to breastfeed. It is not known if Evrysdi passes into breast milk and may harm your baby

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).



Taking Evrysdi: dos and don'ts

DO NOT

- ✗ Your prescription is for Evrysdi tablets, so do not use Evrysdi if it arrives in powder or liquid form. Contact your specialty pharmacy immediately if you receive anything but Evrysdi in tablet form
- ✗ Do not use Evrysdi tablets if the bottle is damaged
- ✗ Do not put the medication in a different container from the one it comes in
- ✗ Do not give Evrysdi tablets with a gastrostomy tube (G-tube) or nasogastric tube (NG-tube)
- ✗ Do not chew, cut, crush, or dissolve the tablet on your tongue
- ✗ Do not mix Evrysdi tablets with food, formula, milk, or with any liquids other than bottled water
- ✗ Do not get the Evrysdi tablet mixture on your skin or in your eyes. If it gets on your skin, wash with soap and water. If it gets in your eyes, flush them with water
- ✗ Do not take the Evrysdi tablet mixture if more than 10 minutes have passed since mixing it with bottled water
- ✗ Do not take an extra dose if you vomit at any time after taking Evrysdi
- ✗ Do not take or give Evrysdi if it has been exposed to temperatures above 86°F (30°C)
- ✗ Do not expose the Evrysdi tablet mixture to sunlight



If you have questions about how to take Evrysdi, contact your healthcare provider. A PAL may also be able to answer questions about Evrysdi; to connect with a PAL, call **(833) 387-9734**, Monday-Friday, 9 AM to 8 PM ET

Important Safety Information (continued)

- **Tell your healthcare provider about all the medicines you take**
- If you were prescribed Evrysdi for Oral Solution, you should receive Evrysdi from the pharmacy as a liquid. If the medicine in the bottle is a powder, **do not use it**. Contact your pharmacist for a replacement
- Avoid getting Evrysdi on your skin or in your eyes. If Evrysdi gets on your skin, wash the area with soap and water. If Evrysdi gets in your eyes, rinse your eyes with water

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).




Support: we're here for you

MySMA Support is a service from Genentech that can help provide information about Evrysdi*

The team is available to:



Answer your questions



Provide information about Evrysdi, including how to take it



Help you understand your insurance coverage



Refer you to appropriate Evrysdi financial assistance options



Coordinate the preparation and delivery of your medicine with the specialty pharmacy



To contact the MySMA Support team: **Visit [Evrysdi.com/PAL](https://evrysdi.com/PAL)** or call **(833) 387-9734** (Monday-Friday, 9 AM to 8 PM ET)

MySMA Support, including the PAL, does not provide medical advice and is not a substitute for your medical team.

Your healthcare provider should always be your main resource for any questions about your health and medical care.

*Enrollment in MySMA Support through the Evrysdi Start Form is mandatory to receive assistance through the program. Participation in MySMA Support is not necessary to receive treatment with Evrysdi.

Important Safety Information (continued)

• **The most common side effects of Evrysdi include:**

- For later-onset SMA: fever, diarrhea, rash
- For infantile-onset SMA: fever; diarrhea; rash; runny nose, sneezing and sore throat (upper respiratory infection); lung infection (lower respiratory infection); constipation; vomiting; cough

These are not all of the possible side effects of Evrysdi. For more information on the risk and benefits profile of Evrysdi, ask your healthcare provider or pharmacist.

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).



Support: we're here for you (continued)

PLANNING AHEAD FOR RENEWALS

If you and your healthcare provider decide to continue with your Evrysdi prescription, your insurance company may require specific tests every 6 to 12 months to approve the renewal of your Evrysdi prescription. It's a good idea to plan ahead for any appointments you may need in advance. The MySMA Support team can help you determine what may be required.

MEET THE TEAM

Your primary contact: **PALs**

Partnership and Access Liaisons (PALs) are the local, main points of contact from Genentech here to support you. Contact a PAL for answers to questions about Evrysdi (including how it works, how to take it, what results were seen in clinical trials, and Important Safety Information), for help understanding your insurance coverage and potential financial support options, and to be connected to helpful resources.

Additional support: **Case Managers**

While a PAL is your main point of contact, Case Managers partner closely with PALs and healthcare providers to help you understand the health insurance process and identify potential financial support options for Evrysdi.

MySMA Support, including the PAL, does not provide medical advice and is not a substitute for your medical team. Your healthcare provider should always be your main resource for any questions about your health and medical care.

Important Safety Information (continued)

- **Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:**
 - are pregnant or plan to become pregnant, as Evrysdi may harm your unborn baby. Ask your healthcare provider for advice before taking this medicine
 - are a woman who can become pregnant:
 - Before you start your treatment with Evrysdi, your healthcare provider may test you for pregnancy
 - Talk to your healthcare provider about birth control methods that may be right for you. Use birth control while on treatment and for at least 1 month after stopping Evrysdi

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).



Support: financial options

Your insurance coverage affects the type of financial support you may be eligible for

Depending on your insurance, support options may include one of the following:



Evrysdi Co-pay Program*

You pay as little as \$0 per prescription for Evrysdi if you qualify for the Evrysdi Co-pay program. The program covers the rest, up to \$25,000 per calendar year.

*This Evrysdi Co-pay Program is valid ONLY for patients with commercial insurance who have a valid prescription for a Food and Drug Administration (FDA)-approved indication of a Genentech medication. Patients using Medicare, Medicaid or any other federal or state government program to pay for their medicines are not eligible. Full program terms and conditions can be found at [EvrysdiCopy.com](https://www.evrysdi.com/copy).



Independent co-pay foundations†

If you need help with the co-pay for your Evrysdi, we can refer you to an independent co-pay assistance foundation.

†Independent co-pay assistance foundations have their own rules for eligibility. Genentech has no involvement or influence in independent foundation decision-making or eligibility criteria and does not know if a foundation will be able to help you. We can only refer you to a foundation that supports your disease state. Genentech does not endorse or show preference for any particular foundation. The foundations we refer you to may not be the only ones that might be able to help you.



The Genentech Patient Foundation†

The Genentech Patient Foundation gives free Evrysdi to people who don't have insurance coverage or who have financial concerns and meet eligibility criteria.

†If you have health insurance, you should try to get other types of financial assistance, if available. You also need to meet income requirements. If you do not have insurance, or if your insurance does not cover your Genentech medicine, you must meet a different set of income requirements. Genentech reserves the right to modify or discontinue the program at any time and to verify the accuracy of information submitted.

For more information

A detailed guide to Insurance and Financial Assistance Options can be found on our website

If you have questions about possible financial support options, contact MySMA Support at **(833) 387-9734** (Monday-Friday, 9 AM to 8 PM ET)

Important Safety Information (continued)

- **Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:**
 - are a woman who can become pregnant:
 - **Pregnancy Registry.** There is a pregnancy registry for women who take Evrysdi during pregnancy. The purpose of this registry is to collect information about the health of the pregnant woman and her baby. If you are pregnant or become pregnant while receiving Evrysdi, tell your healthcare provider right away. Talk to your healthcare provider about registering with the Evrysdi Pregnancy Registry. Your healthcare provider can enroll you in this registry or you can enroll by calling **1-833-760-1098** or visiting **<https://www.evrysdi.com/pregnancyregistry.com>**.

Please see additional **Important Safety Information** throughout and full **Prescribing Information**.



Where To Start

Obtaining Evrysdi

Storing Evrysdi

Taking Evrysdi

Support

FAQs

Important Safety Information

Support: tracking your results

Since everyone’s experience with a new treatment can be different, it can be helpful to keep track of your goals and results over time, and to discuss both with your healthcare provider.

These pages are a simple way to keep your notes organized when starting treatment.

Why I chose Evrysdi: _____

The skills I have now that are important for me to maintain: _____

My expectations for the first 6 months on Evrysdi: _____

My expectations over time (years): _____

How it’s going (how I feel, changes I’ve noticed):

After 1 month _____
After 4 months _____
After 8 months _____
After 1 year _____

My scores (on tests my healthcare provider may use to measure my progress):

Before starting Evrysdi _____
After 6 months _____
After 1 year _____
After 1.5 years _____
After 2 years _____

Important Safety Information (continued)

- **Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:**
 - are an adult male. Evrysdi may affect a man’s ability to have children (fertility). Ask a healthcare provider for advice before taking this medicine
 - are breastfeeding or plan to breastfeed. It is not known if Evrysdi passes into breast milk and may harm your baby



Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).

Support: questions for your healthcare provider

Questions to discuss with my healthcare provider:

- Should I adjust any of the medications I am taking (including vitamins, fiber supplements, laxatives, and prescription medicines)?
- Are there any aspects of my nutrition that we may need to adjust, including my diet or typical meal times? Would it be helpful to talk to a nutritionist?
- What side effects could I experience when starting treatment or as I continue on treatment?
- If I do experience side effects, what should my plan be to address them?
- How should I expect to feel on treatment?
- What results should I expect from treatment?
- How will we measure my progress?
- Are there any tests I may need to plan for to renew my prescription each year?

Additional questions:

This welcome kit includes a **blank journal** you can use to track your results and make notes, and a **reusable cling** where you can write key contact information and keep track of your dosing for easy reference

Important Safety Information (continued)

- **Tell your healthcare provider about all the medicines you take**
- If you were prescribed Evrysdi for Oral Solution, you should receive Evrysdi from the pharmacy as a liquid. If the medicine in the bottle is a powder, **do not use it**. Contact your pharmacist for a replacement
- Avoid getting Evrysdi on your skin or in your eyes. If Evrysdi gets on your skin, wash the area with soap and water. If Evrysdi gets in your eyes, rinse your eyes with water

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).



Support: things you can do about common side effects

Every time you start a new medication, it’s important to know about the potential side effects, what to look out for, and steps to take if they occur.

As always, your healthcare provider is the best resource for any questions. It’s important to talk to your healthcare provider if you notice any changes, begin any new medication, or make changes in your overall care. Below are some tips that may help if you experience these changes.

If you experience diarrhea:

- It means you have loose, watery, and more-frequent bowel movements
- Some dietary tips you can consider include:
 - Drinking more fluids, such as water, broths, and juices
 - Avoiding juices or foods that have a laxative effect, such as prunes or prune juice
 - Eating semisolid and low-fiber food gradually, such as soda crackers, toast, eggs, rice, or chicken
 - Avoiding dairy products, fatty foods, high-fiber foods, or highly seasoned foods for a few days
- In Evrysdi clinical studies, 17% of adults and children taking Evrysdi experienced diarrhea, with a similar frequency in infants. The majority of cases* (8 out of 10) lasted less than 1 week; about 1 out of 10 cases lasted longer than 2 weeks
- Talk to your healthcare provider if you are taking other medications, vitamins, or dietary supplements, including any stool softeners or additional fiber that may impact bowel movements

If you experience a rash:

- It means you have an area of irritated or swollen skin; it may be itchy and red
- Some tips you can consider include:
 - Using moisturizer or lotions
 - Taking a cool bath
 - Avoiding harsh soaps
 - Using cortisone cream to relieve swelling (if not infected)
 - Avoiding wearing tight clothing

*Cases with a known duration.

Important Safety Information (continued)

- **The most common side effects of Evrysdi include:**
 - For later-onset SMA: fever, diarrhea, rash
 - For infantile-onset SMA: fever; diarrhea; rash; runny nose, sneezing and sore throat (upper respiratory infection); lung infection (lower respiratory infection); constipation; vomiting; cough

These are not all of the possible side effects of Evrysdi. For more information on the risk and benefits profile of Evrysdi, ask your healthcare provider or pharmacist.

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).



Frequently asked questions

Obtaining Evrysdi tablets

- ✓ **Can I get Evrysdi tablets from a retail pharmacy?**
Evrysdi tablets are only available through a specialty pharmacy. Once prescribed, they will be shipped directly to your door.
- ✓ **If my delivery arrives later than expected, how do I know it's safe for use?**
Contact your specialty pharmacy representative right away to confirm your product is safe to take.
- ✓ **What should I do if there is an issue with my order, if any of the contents of my shipment (bottle and/or tablets) are damaged, or if the seal on my medication is broken?**
Contact your specialty pharmacy representative right away.

" I love that Evrysdi is an oral treatment I can take at home. Living with a disability is already challenging, and I wanted to find **an option that better fits my lifestyle.**

-Janelle, living with Type 2 SMA **"**



Important Safety Information (continued)

- **Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:**
 - are pregnant or plan to become pregnant, as Evrysdi may harm your unborn baby. Ask your healthcare provider for advice before taking this medicine
 - are a woman who can become pregnant:
 - Before you start your treatment with Evrysdi, your healthcare provider may test you for pregnancy
 - Talk to your healthcare provider about birth control methods that may be right for you. Use birth control while on treatment and for at least 1 month after stopping Evrysdi

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).



Frequently asked questions

Getting started

Where To
Start



What do I need to know about nutrition when starting Evrysdi?

Before starting treatment, tell your healthcare provider what foods you or the person in your care generally eats, as well as your typical meal times. Discuss if any part of your routine may need to be adjusted, such as dairy products, fatty foods, high-fiber foods, highly seasoned foods, or foods that have a laxative effect.

Obtaining
Evrysdi



What side effects should I expect when starting Evrysdi?

Everyone reacts differently when starting a new treatment. Side effects can occur for a number of reasons, including interactions with other medications you may be taking.

Storing
Evrysdi

Before starting treatment, tell your healthcare provider if you or the person in your care is taking any other medications, vitamins, or dietary supplements, including any stool softeners or additional fiber that may impact bowel movements. Your healthcare provider may wish to adjust these medications prior to starting Evrysdi.

Taking
Evrysdi

The most common side effects of Evrysdi for later-onset SMA and infantile-onset SMA include fever, diarrhea, and rash. Infantile-onset SMA also included upper respiratory infection, lower respiratory infection, constipation, vomiting, and cough.

In clinical trials, 17% of adults and children and 15% of infants taking Evrysdi experienced diarrhea. The majority (8 out of 10) of cases of diarrhea lasted less than 1 week and 12% of cases lasted longer than 2 weeks.* See page 21 for some helpful tips if you experience diarrhea, and be sure to speak with your healthcare provider.

Support



What if a dose is missed?

The missed dose should be taken as soon as possible, but no more than 6 hours after the usual dosing time. If more than 6 hours have passed, skip the missed dose altogether and take or give the next dose the following day at your regularly scheduled time.

FAQs



Can I get a prescription for both the liquid and tablet forms of Evrysdi so I can mix and match based on my needs?

Your ability to switch between formulations will depend on discussions with both your healthcare and insurance providers. Be sure to ask your healthcare provider which option is best for you.



Who should I call if I have other questions about Evrysdi?

Your healthcare provider should always be your main resource for any questions about your health and medical care. You can also contact MySMA Support to connect with a PAL, who can answer questions about Evrysdi, and refer you to helpful resources. PALs can also help you understand your insurance and financial support options. Contact MySMA Support at **(833) 387-9734** (Monday-Friday, 9 AM to 8 PM ET).

Important
Safety
Information

*Cases with a known duration.

Important Safety Information (continued)







- **Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:**
 - are a woman who can become pregnant:
 - **Pregnancy Registry.** There is a pregnancy registry for women who take Evrysdi during pregnancy. The purpose of this registry is to collect information about the health of the pregnant woman and her baby. If you are pregnant or become pregnant while receiving Evrysdi, tell your healthcare provider right away. Talk to your healthcare provider about registering with the Evrysdi Pregnancy Registry. Your healthcare provider can enroll you in this registry or you can enroll by calling **1-833-760-1098** or visiting **<https://www.evrysdipregnancyregistry.com>**.

Please see additional **Important Safety Information** throughout and full **Prescribing Information**.



Frequently asked questions

Taking Evrysdi tablets

-  **Why can't I swallow Evrysdi tablets whole with liquids other than water?**
The safety of Evrysdi tablets was studied when swallowed whole with water. The safety of swallowing them any other way is unknown and not recommended. Please read the Dosage and Administration section of the Prescribing Information for more information about taking Evrysdi tablets.
-  **Can the tablets only be swallowed whole?**
Evrysdi tablets can either be swallowed whole with water or mixed with room temperature bottled water. They cannot be chewed, crushed, cut, or dissolved on the tongue.
-  **Why do I need to use room temperature bottled water when preparing the Evrysdi tablet mixture?**
To help make sure every dose is consistent, always mix it with bottled water. Chlorine in tap water can interact with the medicine in Evrysdi.
-  **Can Evrysdi mixture be exposed to sunlight?**
If Evrysdi tablets are taken as a mixture, the mixture must be kept out of sunlight. For example, if sunlight is shining through a window directly on your kitchen table, prepare the mixture someplace else, where the sunlight is indirect and not as strong.
-  **Can I take or give the tablet mixture with a G-tube or NG-tube?**
No, unfortunately it cannot be administered via a gastrostomy tube (G-tube) or nasogastric tube (NG-tube). If you or the person you are caring for has a G-tube or NG-tube, you should ask your healthcare provider about taking the Evrysdi liquid form.
-  **Should Evrysdi tablets be taken with food?**
Evrysdi tablets can be taken with or without food. It will not affect how the tablet works either way. Just remember to take it at the same time each day.

Important Safety Information (continued)

- **Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:**
 - are an adult male. Evrysdi may affect a man's ability to have children (fertility). Ask a healthcare provider for advice before taking this medicine
 - are breastfeeding or plan to breastfeed. It is not known if Evrysdi passes into breast milk and may harm your baby

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).



Frequently asked questions

Ira,
living with
Type 2 SMA

Storage and traveling

- ✓ **I'm switching from Evrysdi liquid to Evrysdi tablets, do I need to take a break in between?**
No need to take a break. If you're switching from Evrysdi liquid to Evrysdi tablets, you can start taking the tablets without missing a dose.
- ✓ **Is there sucralose in Evrysdi tablets?**
No, sucralose is not an ingredient in Evrysdi tablets.
- ✓ **Do Evrysdi tablets have a taste?**
Evrysdi tablets are film coated and do not have a taste when swallowed whole. However, the tablet mixture is strawberry flavored, just like the liquid form of Evrysdi.
- ✓ **What should I do if a dose is vomited, spit up, or not fully swallowed after taking or giving a dose of Evrysdi?**
Do not take or give any more medication that day. Wait until the next day to take or give the next dose at the usual dosing time. If you or the one you care for continues to vomit or spit up while taking Evrysdi, talk to your healthcare provider.



Where To
Start

Obtaining
Evrysdi

Storing
Evrysdi

Taking
Evrysdi

Support

FAQs

Important
Safety
Information

Important Safety Information (continued)

- **Tell your healthcare provider about all the medicines you take**
- If you were prescribed Evrysdi for Oral Solution, you should receive Evrysdi from the pharmacy as a liquid. If the medicine in the bottle is a powder, **do not use it**. Contact your pharmacist for a replacement
- Avoid getting Evrysdi on your skin or in your eyes. If Evrysdi gets on your skin, wash the area with soap and water. If Evrysdi gets in your eyes, rinse your eyes with water

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).



Frequently asked questions

Storage and traveling (continued)

- ✓ **How should I store Evrysdi tablets?**
Evrysdi tablets should be stored at room temperature (68°F to 77°F, 20°C to 25°C).
- ✓ **I've noticed some discoloration/spots on my Evrysdi tablets. Are they still safe and effective?**
If tablets are stored for more than 3 days at elevated humidity and/or temperature (30°C/75% RH), you may notice discoloration or spots on them. Please note you can still use the tablets, as these changes in their appearance have no impact on how Evrysdi works.
- ✓ **Can I travel with Evrysdi tablets?**
Evrysdi tablets can be taken at home or on the go, so you have flexibility in how and where you take them. You can store and take them wherever is most convenient for you, as long as they are kept at room temperature (68°F to 77°F, 20°C to 25°C).
- ✓ **Can I store the tablet in a pill organizer?**
The tablets are sensitive to moisture, so you should not transfer them to a pill organizer. Instead, keep them in the bottle you received them in and always remember to close the bottle. If you've already been storing them in a pill organizer, just transfer them back to the bottle for storage moving forward.

Liz, living
with Type 3
SMA

Important Safety Information (continued)

- **The most common side effects of Evrysdi include:**
 - For later-onset SMA: fever, diarrhea, rash
 - For infantile-onset SMA: fever; diarrhea; rash; runny nose, sneezing and sore throat (upper respiratory infection); lung infection (lower respiratory infection); constipation; vomiting; cough

These are not all of the possible side effects of Evrysdi. For more information on the risk and benefits profile of Evrysdi, ask your healthcare provider or pharmacist.

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).



Important Safety Information

Where To
Start

What is Evrysdi?

Evrysdi is a prescription medicine used to treat spinal muscular atrophy (SMA) in children and adults.

Important Safety Information

Obtaining
Evrysdi

Storing
Evrysdi

Taking
Evrysdi

Support

FAQs

Important
Safety
Information

- **Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:**



- are pregnant or plan to become pregnant, as Evrysdi may harm your unborn baby. Ask your healthcare provider for advice before taking this medicine
- are a woman who can become pregnant:
 - Before you start your treatment with Evrysdi, your healthcare provider may test you for pregnancy
 - Talk to your healthcare provider about birth control methods that may be right for you. Use birth control while on treatment and for at least 1 month after stopping Evrysdi
 - **Pregnancy Registry.** There is a pregnancy registry for women who take Evrysdi during pregnancy. The purpose of this registry is to collect information about the health of the pregnant woman and her baby. If you are pregnant or become pregnant while receiving Evrysdi, tell your healthcare provider right away. Talk to your healthcare provider about registering with the Evrysdi Pregnancy Registry. Your healthcare provider can enroll you in this registry or you can enroll by calling 1-833-760-1098 or visiting <https://www.evrysdipregnancyregistry.com>.



- are an adult male. Evrysdi may affect a man's ability to have children (fertility). Ask a healthcare provider for advice before taking this medicine




- are breastfeeding or plan to breastfeed. It is not known if Evrysdi passes into breast milk and may harm your baby



- **Tell your healthcare provider about all the medicines you take**



- If you were prescribed Evrysdi for Oral Solution, you should receive Evrysdi from the pharmacy as a liquid. If the medicine in the bottle is a powder, **do not use it.** Contact your pharmacist for a replacement



- Avoid getting Evrysdi on your skin or in your eyes. If Evrysdi gets on your skin, wash the area with soap and water. If Evrysdi gets in your eyes, rinse your eyes with water



- **The most common side effects of Evrysdi include:**

- For later-onset SMA: fever, diarrhea, rash
- For infantile-onset SMA: fever; diarrhea; rash; runny nose, sneezing and sore throat (upper respiratory infection); lung infection (lower respiratory infection); constipation; vomiting; cough

These are not all of the possible side effects of Evrysdi. For more information on the risk and benefits profile of Evrysdi, ask your healthcare provider or pharmacist.

You may report side effects to the FDA at **1-800-FDA-1088** or www.fda.gov/medwatch. You may also report side effects to Genentech at **1-888-835-2555**.



Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).

Ira, living with Type 2 SMA



I like taking Evrysdi. It's part of my routine. I don't have to worry about going to the hospital or to a doctor's office to take my dose.

-Janelle,
living with Type 2 SMA

Jim, living with Type 3 SMA



Important Safety Information (continued)

- **Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:**
 - are pregnant or plan to become pregnant, as Evrysdi may harm your unborn baby. Ask your healthcare provider for advice before taking this medicine
 - are a woman who can become pregnant:
 - Before you start your treatment with Evrysdi, your healthcare provider may test you for pregnancy

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#).